

# ALMOST SPANNING SUBGRAPHS OF RANDOM GRAPHS AFTER ADVERSARIAL EDGE REMOVAL

JULIA BÖTTCHER, YOSHIHARU KOHAYAKAWA, AND ANUSCH TARAZ

*Dedicated to Vojtěch Rödl on the occasion of his sixtieth birthday*

**ABSTRACT.** Let  $\Delta \geq 2$  be a fixed integer. We show that the random graph  $\mathcal{G}_{n,p}$  with  $p \geq c(\log n/n)^{1/\Delta}$  is robust with respect to the containment of almost spanning bipartite graphs  $H$  with maximum degree  $\Delta$  and sublinear bandwidth in the following sense. If an adversary deletes arbitrary edges in  $\mathcal{G}_{n,p}$  such that each vertex loses less than half of its neighbours, then asymptotically almost surely the resulting graph still contains a copy of  $H$ .

## 1. INTRODUCTION AND RESULTS

In this paper we study graphs that are robust in the following sense: even after adversarial removal of a specified proportion of their edges, they still contain copies of every graph from a certain class of graphs.

In order to make this precise, we use the notion of *resilience* (see [26]). Let  $\mathcal{P}$  be a monotone increasing graph property and  $G = (V, E)$  be a graph. The *global resilience*  $R_g(G, \mathcal{P})$  of  $G$  with respect to  $\mathcal{P}$  is the minimum  $r \in \mathbb{R}$  such that deleting a suitable set of  $r \cdot |E|$  edges from  $E$  creates a graph which is not in  $\mathcal{P}$ . The *local resilience*  $R_\ell(G, \mathcal{P})$  of  $G$  with respect to  $\mathcal{P}$  is the minimum  $r \in \mathbb{R}$  such that deleting a suitable set of at most  $r \cdot \deg_G(v)$  edges incident to  $v$  for every vertex  $v \in V$  creates a graph which is not in  $\mathcal{P}$ .

For example, using this terminology, the classical theorems of Turán [27] and Dirac [12] can be stated as follows: the global resilience of the complete graph  $K_n$  with respect to containing a clique on  $r$  vertices is  $\frac{1}{r-1} - o(1)$  and the local resilience of  $K_n$  with respect to containing a Hamilton cycle is  $\frac{1}{2} - o(1)$ . In this paper we stay quite close to the scenario of these two examples insofar as we will also consider properties that deal with subgraph containment. However, we are interested in the resilience of graphs which are much sparser than the complete graph.

It turns out that the random graph  $\mathcal{G}_{n,p}$  is well suited for this purpose ( $\mathcal{G}_{n,p}$  is defined on vertex set  $[n] = \{1, \dots, n\}$  and edges exist independently of each other with probability  $p$ ). Sudakov and Vu [26] showed that asymptotically almost surely (a.a.s.) the local resilience of  $\mathcal{G}_{n,p}$  with respect to containing a Hamilton cycle is  $\frac{1}{2} - o(1)$  if  $p > \log^4 n/n$ . A result of Dellamonica et al. [10] implies that a.a.s. the local resilience of  $\mathcal{G}_{n,p}$  with respect to containing cycles of length at least  $(1 - \alpha)n$  is  $\frac{1}{2} - o(1)$  for any  $0 < \alpha < \frac{1}{2}$  and  $p \gg 1/n$ . We shall discuss the various lower bounds for the edge probability  $p$  occurring in these and later results at the end of Section 2.

Recently Balogh, Csaba, and Samotij [7] studied the local resilience of  $\mathcal{G}_{n,p}$  with respect to containing all trees on  $(1 - \eta)n$  vertices with constant maximum degree  $\Delta$ . They showed that there is a constant  $c = c(\Delta, \eta)$  such that for  $p \geq c/n$ , the local resilience of  $\mathcal{G}_{n,p}$  is a.a.s.  $\frac{1}{2} - o(1)$  as well.

Now we extend the scope of investigations to the containment of a much larger class of subgraphs. A graph has *bandwidth* at most  $b$  if there exists a labelling of the vertices by numbers

---

*Date:* March 3, 2010.

The first and third author were partially supported by DFG grant TA 309/2-1. The first author was partially supported by an EUBRANEX grant of the EU programme EM ECW. The second author was partially supported by CNPq (Proc. 308509/2007-2, 485671/2007-7 and 486124/2007-0). The cooperation of the three authors was supported by a joint CAPES-DAAD project (415/ppp-probral/po/D08/11629, Proj. no. 333/09). An extended abstract appeared in: V Latin-American Algorithms, Graphs and Optimization Symposium (LAGOS), Gramado, Brazil, Electronic Notes in Discrete Mathematics, 2009.

$1, \dots, n$ , such that for every edge  $ij$  of the graph we have  $|i - j| \leq b$ . Let  $\mathcal{H}(m, \Delta)$  denote the class of all graphs on  $m$  vertices with maximum degree at most  $\Delta$ , and  $\mathcal{H}_2^b(m, \Delta)$  denote the class of all *bipartite* graphs in  $\mathcal{H}(m, \Delta)$  which have bandwidth at most  $b$ . Our result asserts that the local resilience of  $\mathcal{G}_{n,p}$  with respect to containing any graph  $H$  from  $\mathcal{H}_2^{\beta n}((1 - \eta)n, \Delta)$  is  $\frac{1}{2} - o(1)$  for small  $\beta$  and  $\eta$  and for  $p = p(n) = o(1)$  sufficiently large.

**Theorem 1.** *For each  $\eta, \gamma > 0$  and  $\Delta \geq 2$  there exist positive constants  $\beta$  and  $c$  such that the following holds for  $p \geq c(\log n/n)^{1/\Delta}$ . Asymptotically almost surely every spanning subgraph  $G = (V, E)$  of  $\mathcal{G}_{n,p}$  with  $\deg_G(v) \geq (\frac{1}{2} + \gamma) \deg_{\mathcal{G}_{n,p}}(v)$  for all  $v \in V$  contains a copy of every graph  $H$  in  $\mathcal{H}_2^{\beta n}((1 - \eta)n, \Delta)$ .*

We note that several important classes of graphs have sublinear bandwidth, and hence Theorem 1 does apply to them: this is the case for, e.g., the class of all bounded degree planar graphs (see [8]).

As an application of this theorem we derive a result on polychromatic  $H$ -copies with  $H \in \mathcal{H}_2^{\beta n}((1 - \eta)n, \Delta)$  for certain edge-colourings of  $K_n$  in Section 3. The proof of Theorem 1 is prepared in Sections 4–7 and presented in Section 8. First, however, we will compare our result to related results in the next section.

## 2. BACKGROUND

As we saw at the end of the last section, we are looking for graphs that do not only contain one specific subgraph but a large class of graphs. A graph  $G$  is called *universal* for a class of graphs  $\mathcal{H}$  if  $G$  contains a copy of every graph from  $\mathcal{H}$  as a subgraph. In this section, we first briefly sketch some results concerning universality in general and then come back to resilience with respect to universality.

In [11] it is shown that  $\mathcal{G}_{n,p}$  a.a.s. is universal for  $\mathcal{H}(n, \Delta)$  if  $p = \tilde{\Omega}(n^{-1/2\Delta})$  (where  $\tilde{\Omega}$  hides polylogarithmic factors). It is also shown in [11] that the lower bound for the edge probability  $p$  can be improved if we restrict our attention to balanced bipartite graphs: Let  $\mathcal{H}_2(m, m, \Delta)$  denote the class of bipartite graphs in  $\mathcal{H}(2m, \Delta)$  with two colour classes of equal size. Then  $\mathcal{G}_{2n,p}$  a.a.s. is universal for  $\mathcal{H}_2(n, n, \Delta)$  if  $p = \tilde{\Omega}(n^{-1/\Delta})$ . The same lower bound for  $p$  also guarantees universality for *almost spanning* graphs of arbitrary chromatic number: Alon et al. [4] prove that for every  $\eta > 0$  and  $p = \tilde{\Omega}(n^{-1/\Delta})$ , the random graph  $\mathcal{G}_{n,p}$  a.a.s. is universal for  $\mathcal{H}((1 - \eta)n, \Delta)$ . Alon and Capalbo [2, 3] gave explicit constructions of graphs with average degree  $\tilde{\Omega}(n^{-2/\Delta})n$  that are universal for  $\mathcal{H}(n, \Delta)$ . For results concerning universal graphs for trees see, e.g., [5].

Moving on to resilience, it is clear that an adversary can destroy any spanning subgraph by deleting the edges incident to a single vertex. Hence any graph must have trivial global resilience with respect to universality for spanning subgraphs.

However, if we focus on subgraphs of smaller order, then sparse random graphs have a global resilience arbitrarily close to 1: Alon et al. [4] show that for every  $\gamma > 0$  there is a constant  $\eta > 0$  such that for  $p = \tilde{\Omega}(n^{-1/2\Delta})$  the random graph  $\mathcal{G}_{n,p}$  a.a.s. has global resilience  $1 - \gamma$  with respect to universality for  $\mathcal{H}_2(\eta n, \eta n, \Delta)$ . In other words,  $\mathcal{G}_{n,p}$  contains *many* copies of all graphs from  $\mathcal{H}_2(\eta n, \eta n, \Delta)$  *everywhere*.

Finally, the concept of local resilience allows for non-trivial results concerning universality for almost spanning subgraphs. For example, a conjecture of Bollobás and Komlós proven in [9] asserts that the local resilience of the complete graph  $K_n$  with respect to universality for  $\mathcal{H}_r^{\beta n}(n, \Delta)$  is  $\frac{1}{r} - o(1)$ . Here  $\mathcal{H}_r^{\beta n}(n, \Delta)$  is the class of all  $r$ -colourable  $n$ -vertex graphs with maximum degree at most  $\Delta$  and bandwidth at most  $\beta n$ , and one can show that the bandwidth constraint cannot be omitted.

**Theorem 2.** *For all  $r, \Delta \in \mathbb{N}$  and  $\gamma > 0$ , there exist constants  $\beta > 0$  and  $n_0 \in \mathbb{N}$  such that for every  $n \geq n_0$  the following holds. If  $H$  is an  $r$ -chromatic graph on  $n$  vertices with  $\Delta(H) \leq \Delta$ , and bandwidth at most  $\beta n$  and if  $G$  is a graph on  $n$  vertices with minimum degree  $\delta(G) \geq (\frac{r-1}{r} + \gamma)n$ , then  $G$  contains a copy of  $H$ .*

Our Theorem 1 replaces  $K_n$  by the much sparser graph  $\mathcal{G}_{n,p}$ , but it only treats almost spanning subgraphs and the case  $r = 2$ .

Before we conclude this section, let us briefly explain the lower bounds for the edge probability  $p$  mentioned in the results above, summarized in Table 1.

First, a straightforward counting argument shows that any graph that is universal for  $\mathcal{H}(n, \Delta)$  must have at least  $\Omega(n^{2-2/\Delta})$  edges. Moreover, it is easy to see that an edge probability  $p = n^{\varepsilon-2/\Delta}$  with  $\varepsilon < \frac{1}{\Delta^2}$  is not sufficient to guarantee that  $\mathcal{G}_{n,p}$  is universal for the even more restrictive class  $\mathcal{H}_2(\eta n, \eta n, \Delta)$ . Indeed, consider the graph  $H \in \mathcal{H}_2(\eta n, \eta n, \Delta)$  consisting of  $\eta n/\Delta$  copies of  $K_{\Delta, \Delta}$ . The expected number of copies of  $K_{\Delta, \Delta}$  in  $\mathcal{G}_{n,p}$  is at most

$$n^{2\Delta} p^{\Delta^2} = n^{2\Delta} (n^{-\frac{2}{\Delta} + \varepsilon})^{\Delta^2} = n^{2\Delta - 2\Delta + \varepsilon \Delta^2} \ll n,$$

and hence a.a.s.  $\mathcal{G}_{n,p}$  does not contain a copy of  $H$ .

|              | Result                                                                                            | $p$                  | Reference |
|--------------|---------------------------------------------------------------------------------------------------|----------------------|-----------|
| Universality | $\mathcal{H}_2(n, n, \Delta) \subseteq \mathcal{G}_{2n,p}$                                        | $p = n^{-1/\Delta}$  | [11]      |
|              | $\mathcal{H}(n, \Delta) \subseteq \mathcal{G}_{n,p}$                                              | $p = n^{-1/2\Delta}$ | [11]      |
|              | $\mathcal{H}((1-\eta)n, \Delta) \subseteq \mathcal{G}_{n,p}$                                      | $p = n^{-1/\Delta}$  | [4]       |
| Resilience   | $R_g(\mathcal{G}_{n,p}, \mathcal{H}_2(\eta n, \eta n, \Delta)) \geq 1 - \gamma$                   | $p = n^{-1/2\Delta}$ | [10]      |
|              | $R_\ell(\mathcal{G}_{n,p}, \mathcal{H}_2^{\beta n}((1-\eta)n, \Delta)) \geq \frac{1}{2} - \gamma$ | $p = n^{-1/\Delta}$  | Theorem 1 |

TABLE 1. Summary of (best) known universality and resilience results (logarithmic factors for  $p$  are omitted).

### 3. AN APPLICATION: POLYCHROMATIC COPIES OF BIPARTITE GRAPHS

Let  $\varphi$  be an arbitrary colouring of the edges of the complete graph  $K_n$ . If  $\varphi$  uses no colour more than  $k$  times then we say that  $\varphi$  is  $k$ -bounded. Moreover, a copy of a graph  $H$  in  $K_n$  is *polychromatic* if  $\varphi$  uses no colour more than once on  $H$ . If there is a polychromatic copy of  $H$  in  $K_n$  then  $\varphi$  is called  $H$ -polychromatic.

Erdős, Nešetřil, and Rödl [13] asked for which  $k = k(n)$  every  $k$ -bounded edge colouring of  $K_n$  has a polychromatic Hamilton cycle. Frieze and Reed [14] showed that  $k(n)$  can grow as fast as  $\kappa n / \log n$  for some constant  $\kappa$  (for early progress on this problem see the references in [14]). Albert, Frieze, and Reed [1] improved this bound to  $n/65$ , which shows that  $k$  can grow linearly, as was previously conjectured by Hahn and Thomassen [17].

Here we consider the analogous question for  $H$ -polychromatic colourings with  $H \in \mathcal{H}_2^{\beta n}((1-\eta)n, \Delta)$ . As a consequence of our main theorem, Theorem 1, we prove the following result.

**Theorem 3.** *For every  $\eta > 0$  and  $\Delta \geq 2$  there exist positive constants  $\beta$  and  $\kappa$  such that for  $n$  sufficiently large, for every graph  $H \in \mathcal{H}_2^{\beta n}((1-\eta)n, \Delta)$  and  $k \leq \kappa(n/\log n)^{1/\Delta}$ , every  $k$ -bounded edge-colouring of  $K_n$  is  $H$ -polychromatic.*

For the proof of this theorem we apply the strategy of [14] and do the following for a given  $k$ -bounded edge colouring  $\varphi$  of  $K_n$ . We first take a random subgraph  $\Gamma = \mathcal{G}_{n,p}$  of  $K_n$  and then delete all edges in  $\Gamma$  whose colour appears more than once in  $\Gamma$ . Denote the resulting graph by  $\Gamma(\varphi)$ . Any subgraph of  $\Gamma(\varphi)$  is trivially polychromatic and hence it remains to show that there is a copy of  $H$  in  $\Gamma(\varphi)$  in order to establish Theorem 3. In view of Theorem 1 it clearly suffices to prove the following lemma.

**Lemma 4.** *Let  $p = p(n)$  and  $k = k(n)$  be such that  $p \geq 10^6 \log n/n$  and  $pk \leq 10^{-3}$ . For any  $k$ -bounded edge colouring  $\varphi$  of  $K_n$ , with probability  $1 - o(1)$  all vertices  $v$  in  $\Gamma = \mathcal{G}_{n,p}$  satisfy  $\deg_{\Gamma(\varphi)}(v) \geq \frac{2}{3} \deg_{\Gamma}(v)$ .*

*Proof (sketch).* Let  $v$  be an arbitrary vertex of  $\Gamma$ . We distinguish two different types of edges in  $E(v, N_\Gamma(v) \setminus N_{\Gamma(\varphi)}(v))$ : The set  $N_1$  of those edges whose colour appears only once in  $N_\Gamma(v)$  (but also somewhere else in  $\Gamma$ ) and the set  $N_2$  of those edges whose colour appears at least twice in  $N_\Gamma(v)$ . With probability  $1 - o(1/n)$  we have that  $\deg_\Gamma(v)$  lies in the interval  $[(1 - \frac{1}{20})np, (1 + \frac{1}{20})np]$ . Therefore, showing

- (i)  $\mathbb{P}(|N_1| \geq \frac{1}{10}np) = o(1/n)$  and
- (ii)  $\mathbb{P}(|N_2| \geq \frac{1}{10}np) = o(1/n)$

and applying the union bound proves the lemma.

For establishing (i) we expose the edges incident to  $v$  first, which enables us to determine  $\deg_\Gamma(v)$ . We have  $\mathbb{P}(\deg_\Gamma(v) \geq \frac{21}{20}np) \leq o(1/n)$ . Subsequently we expose the remaining edges. Recall that for any edge  $vw \in N_1$  the colour  $\varphi(vw)$  appears somewhere else in  $\Gamma$ , which happens with probability at most  $p' := pk$ . Since these events are independent for different colours, we have

$$\begin{aligned} \mathbb{P}(|N_1| \geq t) &\leq \mathbb{P}(\deg_\Gamma(v) \geq \frac{21}{20}np) + \mathbb{P}(|N_1| > t \mid \deg_\Gamma(v) \leq \frac{21}{20}np) \\ &\leq o(1/n) + \mathbb{P}(X \geq t), \end{aligned}$$

where  $X$  is a random variable with distribution  $\text{Bi}(n', p')$  where  $n' = \deg_\Gamma(v) \leq \frac{21}{20}np$ . Clearly  $\mathbb{E} X \leq \frac{21}{20}np \cdot pk \leq \frac{1}{100}np$  and therefore (i) follows from two applications of a Chernoff type bound since  $np \geq 10^6 \log n$ .

For establishing (ii) consider the random variable  $Y$  that counts edges in  $N_\Gamma(v)$  whose colour appears only once in  $N_\Gamma(v)$ . Then  $|N_2| = \deg_\Gamma(v) - Y$  and so it suffices to show that  $\mathbb{P}(Y \leq \frac{19}{20}np) = o(1/n)$ , using again that  $\deg_\Gamma(v) > \frac{21}{20}np$  happens with probability  $o(1/n)$ . To see this, assume that  $1, \dots, \ell$  are the colours that appear on the edges of  $K_n$  containing  $v$ , and let  $k_i$  be the number of such edges with colour  $i \in [\ell]$ . Then  $Y = \sum_{i \in [\ell]} Y_i$  where  $Y_i$  is the indicator variable for the event that  $N_\Gamma(v)$  contains exactly one edge of colour  $i$ . Observe that the  $Y_i$  are independent random variables and that  $\mathbb{P}(Y_i = 1) = k_i p (1 - p)^{k_i - 1}$ . In addition  $1 \geq (1 - p)^{k_i - 1} \geq (1 - p)^k \geq \exp(-\frac{p}{1-p}k) \geq \frac{100}{101}$  and hence

$$\mathbb{E} Y = \sum_{i \in [\ell]} k_i p (1 - p)^{k_i - 1} \leq np \quad \text{and} \quad \mathbb{E} Y \geq \frac{100}{101}(n - 1)p \geq \frac{99}{100}np.$$

We conclude  $\mathbb{P}(Y \leq \frac{19}{20}np) = o(1/n)$  from  $\mathbb{P}(Y \leq \mathbb{E} Y - t) \leq \exp(-\frac{1}{2}t^2/\mathbb{E} Y)$  (see [19, Theorem 2.10]) by setting  $t := \frac{1}{100}np$  and using  $np \geq 10^6 \log n$ .  $\square$

As explained earlier the bound on  $k(n)$  established in [14] is not best possible. We also believe that the bound on  $k(n)$  in Theorem 3 can be improved. In fact, the only upper bound we are aware of is the trivial bound  $k(n) < n/\Delta$ .

#### 4. SPARSE REGULARITY

In this section we will introduce one of the basic tools for our proof, a sparse version of the regularity lemma (see [20, 22] and [16]). Before stating this lemma we introduce the necessary definitions.

Let  $G = (V, E)$  be a graph,  $p \in (0, 1]$ , and  $\varepsilon, d > 0$  be reals. For disjoint nonempty  $U, W \subseteq V$  the  $p$ -density of the pair  $(U, W)$  is defined by  $d_{G,p}(U, W) := e_G(U, W)/(p|U||W|)$ .

The pair  $(U, W)$  is  $(\varepsilon, d, p)$ -dense if  $d_{G,p}(U', W') \geq d - \varepsilon$  for all  $U' \subseteq U$  and  $W' \subseteq W$  with  $|U'| \geq \varepsilon|U|$  and  $|W'| \geq \varepsilon|W|$ . Omitting the parameters  $d$ , or  $\varepsilon$  and  $d$ , we may also speak of  $(\varepsilon, p)$ -dense pairs, or  $p$ -dense pairs.

An  $(\varepsilon, p)$ -dense partition of  $G = (V, E)$  is a partition  $V_0 \dot{\cup} V_1 \dot{\cup} \dots \dot{\cup} V_r$  of  $V$  with  $|V_0| \leq \varepsilon|V|$  such that  $(V_i, V_j)$  is an  $(\varepsilon, p)$ -dense pair in  $G$  for all but at most  $\varepsilon \binom{r}{2}$  pairs  $ij \in \binom{[r]}{2}$ . The partition classes  $V_i$  with  $i \in [r]$  are called the *clusters* of the partition and  $V_0$  is the *exceptional set*. An  $(\varepsilon, p)$ -dense partition  $V_0 \dot{\cup} V_1 \dot{\cup} \dots \dot{\cup} V_r$  is an  $(\varepsilon, d, p)$ -dense partition of  $G$  with *reduced graph*  $R = ([r], E(R))$  if the pair  $(V_i, V_j)$  is  $(\varepsilon, d, p)$ -dense in  $G$  whenever  $ij \in E(R)$ .

The sparse regularity lemma asserts  $p$ -dense partitions for sparse graphs  $G$  without dense spots. To quantify this latter property we need the following notion. Let  $\eta > 0$  be a real number and

$K > 1$  an integer. We say that  $G = (V, E)$  is  $(\eta, K)$ -bounded with respect to  $p$  if for all disjoint sets  $X, Y \subseteq V$  with  $|X|, |Y| \geq \eta|V|$  we have  $e_G(X, Y) \leq Kp|X||Y|$ .

**Lemma 5** (sparse regularity lemma). *For each  $\varepsilon > 0$ , each  $K > 1$ , and each  $r_0 \geq 1$  there are constants  $r_1, \nu$ , and  $n_0$  such that for any  $p \in (0, 1]$  the following holds. Any graph  $G = (V, E)$  which has at least  $n_0$  vertices and is  $(\nu, K)$ -bounded with respect to  $p$  admits an  $(\varepsilon, p)$ -dense equipartition with  $r$  clusters for some  $r_0 \leq r \leq r_1$ .  $\square$*

It follows directly from the definition that sub-pairs of  $p$ -dense pairs again form  $p$ -dense pairs.

**Proposition 6.** *Let  $(X, Y)$  be  $(\varepsilon, d, p)$ -dense,  $X' \subseteq X$  with  $|X'| = \mu|X|$ . Then  $(X', Y)$  is  $(\frac{\varepsilon}{\mu}, d, p)$ -dense.  $\square$*

In addition neighbourhoods of most vertices in a  $p$ -dense pair are not much smaller than expected. Again, this is a direct consequence of the definition of  $p$ -dense pairs.

**Proposition 7.** *Let  $(X, Y)$  be  $(\varepsilon, d, p)$ -dense. Then less than  $\varepsilon|X|$  vertices  $x \in X$  have  $|N_Y(x)| < (d - \varepsilon)p|Y|$ .  $\square$*

Some properties of the graph  $G$  translate to certain properties of the reduced graph  $R$  of the partition constructed by the sparse regularity lemma. An example of this phenomenon is given in the following lemma, Lemma 8, which is a minimum degree version of the sparse regularity lemma.

**Lemma 8** (sparse regularity lemma, minimum degree version for  $\mathcal{G}_{n,p}$ ). *For all  $\alpha \in [0, 1]$ ,  $\varepsilon > 0$ , and every integer  $r_0$ , there is an integer  $r_1 \geq 1$  such that for all  $d \in [0, 1]$  the following holds a.a.s. for  $\Gamma = \mathcal{G}_{n,p}$  if  $\log^4 n/(pn) = o(1)$ . Let  $G = (V, E)$  be a spanning subgraph of  $\Gamma$  with  $\deg_G(v) \geq \alpha \deg_\Gamma(v)$  for all  $v \in V$ . Then there is an  $(\varepsilon, d, p)$ -dense partition of  $G$  with reduced graph  $R$  of minimum degree  $\delta(R) \geq (\alpha - d - \varepsilon)|V(R)|$  with  $r_0 \leq |V(R)| \leq r_1$ .*

Before we show how Lemma 8 can be deduced from Lemma 5, we remark that we do observe “more” than a mere inheritance of properties here: the graph  $G$  we started with is *sparse*, but the reduced graph  $R$  we obtain in Lemma 8 is *dense*. This will enable us to apply results obtained for dense graphs to the reduced graph  $R$ , and hence use such dense results to draw conclusions about sparse graphs. For the proof we also need the following lemma which collects some well known facts about the edge distribution in random graphs  $\mathcal{G}_{n,p}$  and follows directly from the Chernoff bound for binomially distributed random variables.

**Lemma 9.** *If  $\log^4 n/(pn) = o(1)$  then a.a.s. the random graph  $\Gamma = \mathcal{G}_{n,p}$  has the following properties. For all vertex sets  $X, Y, Z \subseteq V(\Gamma)$  with  $X \cap Y = \emptyset$  and  $|X|, |Y|, |Z| \geq \frac{n}{\log n}$ ,  $|Z| \leq n - \frac{n}{\log n}$  we have*

- (i)  $e_\Gamma(X) = (1 \pm \frac{1}{\log n})p \binom{|X|}{2}$ ,
- (ii)  $e_\Gamma(X, Y) = (1 \pm \frac{1}{\log n})p|X||Y|$ ,
- (iii)  $\sum_{z \in Z} \deg_\Gamma(z) = (1 \pm \frac{1}{\log n})p|Z|n$ .  $\square$

*Proof of Lemma 8.* For the proof we will use the sparse regularity lemma (Lemma 5) and the facts about the edge distribution in random graphs provided by Lemma 9.

Given  $\alpha, \varepsilon$ , and  $r_0$  let  $r_1, \nu$ , and  $n_0$  be as provided by Lemma 5 for input

$$\varepsilon' := \varepsilon^2/100, \quad K := 1 + \varepsilon', \quad \text{and} \quad r'_0 := \min\{2r_0, \lceil 1/\varepsilon' \rceil\}.$$

Let further  $d$  be given and assume that  $n$  is such that  $n \geq n_0$ ,  $\log n \geq 1/\varepsilon'$ , and  $\log n \geq 1/\nu$ . Let  $\Gamma$  be a typical graph from  $\mathcal{G}_{n,p}$  with  $\log^4 n/(pn) = o(1)$ , i.e., a graph satisfying properties (i)–(iii) of Lemma 9. We will show that then  $\Gamma$  also satisfies the conclusion of Lemma 8.

To this end we consider an arbitrary subgraph  $G = (V, E)$  of  $\Gamma$  that satisfies the assumptions of Lemma 8. By property (ii) of Lemma 9 the graph  $G \subseteq \Gamma$  is  $(1/\log n, 1 + 1/\log n)$ -bounded with respect to  $p$ . Since we have  $1 + 1/\log n \leq 1 + \varepsilon' = K$ , the sparse regularity lemma (Lemma 5) with input  $\varepsilon'$ ,  $K$ , and  $r'_0$  asserts that  $G$  has an  $(\varepsilon', p)$ -dense equipartition  $V = V'_0 \dot{\cup} V'_1 \dot{\cup} \dots \dot{\cup} V'_{r'}$  for some  $r'_0 \leq r' \leq r_1$ . Observe that there are at most  $r'\sqrt{\varepsilon'}$  clusters in this partition which are

contained in more than  $r'\sqrt{\varepsilon'}$  pairs that are not  $(\varepsilon', p)$ -dense. We add all these clusters to  $V'_0$ , denote the resulting set by  $V_0$  and the remaining clusters by  $V_1, \dots, V_r$ . Then  $r_0 \leq r'/2 \leq r \leq r_1$  and we claim that the partition  $V = V_0 \dot{\cup} V_1 \dot{\cup} \dots \dot{\cup} V_r$  has the desired properties.

Indeed,  $|V_0| \leq \varepsilon'n + r'\sqrt{\varepsilon'}(n/r') \leq \varepsilon n$  and the number of pairs in  $V_1 \dot{\cup} \dots \dot{\cup} V_r$  which are not  $(\varepsilon, p)$ -dense is at most  $r \cdot r'\sqrt{\varepsilon'} \leq 2r^2\sqrt{\varepsilon'} \leq \varepsilon \binom{r}{2}$ . It follows that  $V_1 \dot{\cup} \dots \dot{\cup} V_r$  is an  $(\varepsilon, p)$ -dense partition and hence an  $(\varepsilon, d, p)$ -dense partition. Let  $R$  be the (edge maximal) corresponding reduced graph, i.e.,  $R$  has vertex set  $[r]$  and edges  $ij$  for exactly all  $(\varepsilon, d, p)$ -dense pairs  $(V_i, V_j)$  with  $i, j \in [r]$ . It remains to show that we have  $\delta(R) \geq (\alpha - d - \varepsilon)|R|$ .

To see this, define  $L := |V_i|$  for all  $i \in [r]$  and consider arbitrary disjoint sets  $X, Y \subseteq V(G)$ . Then  $\sum_{x \in X} \deg_G(x) = 2e_G(X) + e_G(X, Y) + e_G(X, V \setminus (X \cup Y))$  and therefore

$$e_G(X, Y) \geq \left( \alpha \sum_{x \in X} \deg_\Gamma(x) \right) - 2e_\Gamma(X) - e_\Gamma(X, V \setminus (X \cup Y)).$$

By properties (i)–(iii) of Lemma 9 this implies

$$\begin{aligned} e_G(X, Y) &\geq \alpha \left( 1 - \frac{1}{\log n} \right) p |X| n - 2 \left( 1 + \frac{1}{\log n} \right) p \binom{|X|}{2} \\ &\quad - \left( 1 + \frac{1}{\log n} \right) p |X| (n - |X| - |Y|) \\ &\geq (\alpha(1 - \varepsilon')n - (1 + \varepsilon')(n - |Y|)) p |X|, \end{aligned} \tag{1}$$

as long as  $|X| \geq n/\log n$  and  $|X \cup Y| \leq n - n/\log n$ . Now fix  $i \in [r]$  and let  $\bar{V}_i := V \setminus (V_0 \cup V_i)$ . Then

$$e_G(V_i, \bar{V}_i) \leq (\deg_R(i) + 2r\sqrt{\varepsilon'}) (1 + \varepsilon') p L^2 + (r - \deg_R(i)) d p L^2$$

since each cluster is contained in at most  $r'\sqrt{\varepsilon'} \leq 2r\sqrt{\varepsilon'}$  irregular pairs and because  $R$  is an  $(\varepsilon', d, p)$ -reduced graph and  $G \subseteq \Gamma$  is  $(1/\log n, 1 + \varepsilon')$ -bounded with respect to  $p$ . On the other hand, (1) implies that

$$\begin{aligned} e_G(V_i, \bar{V}_i) &\geq \left( \alpha(1 - \varepsilon')n - (1 + \varepsilon')(|V_0| + |V_i|) \right) p |V_i| \\ &\geq \left( \alpha(1 - \varepsilon') - (1 + \varepsilon')3\sqrt{\varepsilon'} \right) p L n \end{aligned}$$

where we used  $|V_0| \leq (\varepsilon' + \sqrt{\varepsilon'})n$  and  $|V_i| \leq n/r'_0 \leq \varepsilon'n$ . We conclude that

$$(\deg_R(i)(1 + \varepsilon' - d) + 2r\sqrt{\varepsilon'}(1 + \varepsilon') + rd) p L^2 \geq (\alpha(1 - \varepsilon') - (1 + \varepsilon')3\sqrt{\varepsilon'}) p r L^2$$

since  $n/L \geq r$ . This gives

$$\begin{aligned} \deg_R(i) &\geq \deg_R(i)(1 + \varepsilon' - d) \geq \left( \alpha(1 - \varepsilon') - (1 + \varepsilon')3\sqrt{\varepsilon'} - 2\sqrt{\varepsilon'}(1 + \varepsilon') - d \right) r \\ &\geq \left( \alpha - \alpha\varepsilon' - 9\sqrt{\varepsilon'} - d \right) r \geq (\alpha - d - \varepsilon)|R|. \end{aligned}$$


Hence the  $(\varepsilon, d, p)$ -dense partition  $V = V_0 \dot{\cup} V_1 \dot{\cup} \dots \dot{\cup} V_r$  has a reduced graph  $R$  with  $\delta(R) \geq (\alpha - d - \varepsilon)|R|$ .  $\square$

## 5. MAIN LEMMAS

In this section we will formulate the main lemmas and outline how they will be combined in Section 8 to give the proof of Theorem 1. For this we first need to define two (families of) special graphs.

For  $r, t \in \mathbb{N}$ ,  $t$  even, let  $U = \{u_1, \dots, u_r\}$ ,  $V = \{v_1, \dots, v_r\}$ ,  $C = \{c_{i,j}, c'_{i,j} : i \in [r], j \in [2t]\}$ , and  $B = \{b_{i,j}, b'_{i,j} : i \in [r], j \in [2t]\}$ . Let the *ladder*  $R_r^* = (U \dot{\cup} V, E(R_r^*))$  have edges  $E(R_r^*) := \{u_i v_j : i, j \in [r], |i - j| \leq 1\}$  and let the *spin graph*  $R_{r,t} = (U \dot{\cup} V \dot{\cup} C \dot{\cup} B, E(R_{r,t}))$  be the graph



FIGURE 1. The ladder  $R_r^*$  and the spin graph  $R_{r,t}$  for the special case  $t = 2$ .

with the following edge set (see Figure 1):

$$E(R_{r,t}) := \bigcup_{\substack{i,i' \in [r], i' \neq i \\ j,j' \in [2t] \\ k,k' \in [t] \\ \ell,\ell' \in [t+1, 2t]}} \left( \left\{ u_i v_i, b_{i,k} b'_{i,k'}, b_{i,\ell} b'_{i,\ell'}, c_{i,k} c'_{i,k'}, c_{i,\ell} c'_{i,\ell'} \right\} \cup \left\{ b_{i,j} v_i, c_{i,j} v_i \right\} \cup \left\{ b'_{i,k} b'_{i,\ell}, c_{i'-1,\ell} c'_{i',k}, c_{i'-1,\ell} c'_{i',k} \right\} \right).$$

Now we can state our four main lemmas, two partition lemmas and two embedding lemmas. We start with the lemma for  $G$ , which constructs a partition of the host graph  $G$ . This lemma is a consequence of the sparse regularity lemma (Lemma 8) and asserts a  $p$ -dense partition of  $G$  such that its reduced graph contains a spin graph. We will indicate below why this is useful for the embedding of  $H$ . The lemma for  $G$  produces clusters of very different sizes: A set of larger clusters  $U_i$  and  $V_i$  which we call *big clusters* and which will accommodate most of the vertices of  $H$  later, and a set of smaller clusters  $B_{i,j}, B'_{i,j}, C_{i,j}$ , and  $C'_{i,j}$ . The  $B_{i,j}$  and  $B'_{i,j}$  are called *balancing clusters* and the  $C_{i,j}$  and  $C'_{i,j}$  *connecting clusters*. They will be used to host a small number of vertices of  $H$ . These vertices balance and connect the pieces of  $H$  that are embedded into the big clusters. The proof of Lemma 10 is given in Section 9.

In the formulation of this lemma (and also in the lemma for  $H$  below) we abuse the notation in the following sense. For two sets  $A$  and  $B$  and a number  $x$  we write  $|A| := |B| \geq x$  by which we simultaneously mean that  $A$  is defined to be the set  $B$  and that the size  $|A| = |B|$  of this set is at least  $x$ .

**Lemma 10** (Lemma for  $G$ ). *For all integers  $t, r_0 > 0$  and reals  $\eta, \gamma > 0$  there are positive reals  $\eta'$  and  $d$  such that for all  $\varepsilon > 0$  there is  $r_1$  such that the following holds a.a.s. for  $\Gamma = \mathcal{G}_{n,p}$  with  $\log^4 n/(pn) = o(1)$ . Let  $G = (V, E)$  be a spanning subgraph of  $\Gamma$  with  $\deg_G(v) \geq (\frac{1}{2} + \gamma) \deg_\Gamma(v)$  for all  $v \in V$ . Then there is  $r_0 \leq r \leq r_1$ , a subset  $V_0$  of  $V$  with  $|V_0| \leq \varepsilon n$ , and a mapping  $g$  from  $V \setminus V_0$  to the spin graph  $R_{r,t}$  such that for every  $i \in [r], j \in [2t]$  we have*

- (G1)  $|U_i| := |g^{-1}(u_i)| \geq (1 - \eta) \frac{n}{2r}$  and  $|V_i| := |g^{-1}(v_i)| \geq (1 - \eta) \frac{n}{2r}$ ,
- (G2)  $|C_{i,j}| := |g^{-1}(c_{i,j})| \geq \eta' \frac{n}{2r}$  and  $|C'_{i,j}| := |g^{-1}(c'_{i,j})| \geq \eta' \frac{n}{2r}$ ,  
 $|B_{i,j}| := |g^{-1}(b_{i,j})| \geq \eta' \frac{n}{2r}$  and  $|B'_{i,j}| := |g^{-1}(b'_{i,j})| \geq \eta' \frac{n}{2r}$ ,
- (G3) the pair  $(g^{-1}(x), g^{-1}(y))$  is  $(\varepsilon, d, p)$ -dense for all  $xy \in E(R_{r,t})$ .

Our second lemma provides a partition of  $H$  that fits the structure of the partition of  $G$  generated by Lemma 10. We will first state this lemma and then explain the different properties which it guarantees.

**Lemma 11** (Lemma for  $H$ ). *For all integers  $\Delta$  there is an integer  $t > 0$  such that for any  $\eta > 0$  and any integer  $r \geq 1$  there is  $\beta > 0$  such that the following holds for all integers  $m$  and all bipartite graphs  $H$  on  $m$  vertices with  $\Delta(H) \leq \Delta$  and  $\text{bw}(H) \leq \beta m$ . There is a homomorphism  $h$  from  $H$  to the spin graph  $R_{r,t}$  such that for every  $i \in [r], j \in [2t]$*

- (H1)  $|\tilde{U}_i| := |h^{-1}(u_i)| \leq (1 + \eta) \frac{m}{2r}$  and  $|\tilde{V}_i| := |h^{-1}(v_i)| \leq (1 + \eta) \frac{m}{2r}$ ,
- (H2)  $|\tilde{C}_{i,j}| := |h^{-1}(c_{i,j})| \leq \eta \frac{m}{2r}$  and  $|\tilde{C}'_{i,j}| := |h^{-1}(c'_{i,j})| \leq \eta \frac{m}{2r}$ ,  
 $|\tilde{B}_{i,j}| := |h^{-1}(b_{i,k})| \leq \eta \frac{m}{2r}$  and  $|\tilde{B}'_{i,j}| := |h^{-1}(b'_{i,k})| \leq \eta \frac{m}{2r}$ ,
- (H3)  $\tilde{C}_{i,j}$ ,  $\tilde{C}'_{i,j}$ ,  $\tilde{B}_{i,j}$ , and  $\tilde{B}'_{i,j}$  are 3-independent in  $H$ ,
- (H4)  $\deg_{\tilde{V}_i}(y) = \deg_{\tilde{V}_i}(y') \leq \Delta - 1$  for all  $yy' \in \binom{\tilde{C}_{i,j}}{2} \cup \binom{\tilde{B}_{i,j}}{2}$ ,  
 $\deg_{\tilde{C}_i}(y) = \deg_{\tilde{C}_i}(y')$  for all  $y, y' \in \tilde{C}'_{i,j}$ ,  
 $\deg_{L(i,j)}(y) = \deg_{L(i,j)}(y')$  for all  $y, y' \in \tilde{B}'_{i,j}$ ,

where  $\tilde{C}_i := \bigcup_{k \in [2t]} \tilde{C}_{i,k}$  and  $L(i, j) := \bigcup_{k \in [2t]} \tilde{B}_{i,k} \cup \bigcup_{k < j} \tilde{B}'_{i,k}$ . Further, let  $\tilde{X}_i$  with  $i \in [r]$  be the set of vertices in  $\tilde{V}_i$  with neighbours outside  $\tilde{U}_i$ . Then

- (H5)  $|\tilde{X}_i| \leq \eta |\tilde{V}_i|$ .

This lemma asserts a homomorphism  $h$  from  $H$  to a spin graph  $R_{r,t}$ . Recall that  $R_{r,t}$  is contained in the reduced graph of the  $p$ -dense partition provided by Lemma 10. As we will see, we can fix the parameters in this lemma such that, when we apply it together with Lemma 10, the homomorphism  $h$  has the following additional property. The number  $\tilde{L}$  of vertices that it maps to a vertex  $a$  of the spin graph is less than the number  $L$  contained in the corresponding cluster  $A$  provided by Lemma 10 (compare (G1) and (G2) with (H1) and (H2) and note that  $m \leq (1 - \eta)n$ ). If  $A$  is a big cluster, then the numbers  $L$  and  $\tilde{L}$  differ only slightly (these vertices will be embedded using the constrained blow-up lemma), but for balancing and connecting clusters  $A$  the number  $\tilde{L}$  is much smaller than  $L$  (this is necessary for the embedding of these vertices using the connection lemma). With property (H5) Lemma 11 further guarantees that only few edges of  $H$  are not assigned either to two connecting or balancing clusters, or to two big clusters. This is helpful because it implies that we do not have to take care of “too many dependencies” between the applications of the blow-up lemma and the connection lemma. The remaining properties (H3)–(H4) of Lemma 11 are technical but required for the application of the connection lemma (see conditions (B) and (C) of Lemma 13).

The vertices in  $\tilde{C}_{i,j}$  and  $\tilde{C}'_{i,j}$  are also called *connecting vertices* of  $H$ , the vertices in  $\tilde{B}_{i,j}$  and  $\tilde{B}'_{i,j}$  *balancing vertices*.

We next describe the two embedding lemmas, the constrained blow-up lemma (Lemma 12) and the connection lemma (Lemma 13), which we would like to use on the partitions of  $G$  and  $H$  provided by Lemmas 10 and 11. The connecting lemma will be used to embed the connecting and balancing vertices into the connecting and balancing clusters *after* all the other vertices are embedded into the big clusters with the help of the constrained blow-up lemma.

The constrained blow-up lemma states that bipartite graphs  $H$  with bounded maximum degree can be embedded into a  $p$ -dense pair  $G = (U, V)$  whose cluster sizes are just slightly bigger than the partition classes of  $H$ . This lemma further guarantees the following. If we specify a small family of small special sets in one of the partition classes of  $H$  and a small family of small forbidden sets in the corresponding cluster of  $G$ , then no special set is mapped to a forbidden set.

The existence of these forbidden sets is in fact a main difference to the classical blow-up lemma which is used in the dense setting, where a small family of special vertices of  $H$  can be guaranteed to be mapped to a *required set* of linear size in  $G$ . This is very useful in a dense graph, because its neighbourhoods (into which we would like to embed neighbours of already embedded vertices) are of linear size. In contrast, the property of having forbidden sets will be crucial for the sparse setting when we will apply this lemma together with the connection lemma in the proof of Theorem 1 in order to handle the “dependencies” between these applications. The proof of this lemma is given in Section 11 and relies on techniques developed in [4].



**Lemma 12** (Constrained blow-up lemma). *For every integer  $\Delta > 1$  and for all positive reals  $d$ , and  $\eta$  there exist positive constants  $\varepsilon$  and  $\mu$  such that for all positive integers  $r_1$  there is  $c$  such that for all integers  $1 \leq r \leq r_1$  the following holds a.a.s. for  $\Gamma = \mathcal{G}_{n,p}$  with  $p \geq c(\log n/n)^{1/\Delta}$ . Let  $G = (U, V) \subseteq \Gamma$  be an  $(\varepsilon, d, p)$ -dense pair with  $|U|, |V| \geq n/r$  and let  $H$  be a bipartite graph on vertex classes  $\tilde{U} \dot{\cup} \tilde{V}$  of sizes  $|\tilde{U}|, |\tilde{V}| \leq (1 - \eta)n/r$  and with  $\Delta(H) \leq \Delta$ . Moreover, suppose that there is a family  $\mathcal{H} \subseteq \binom{\tilde{V}}{\Delta}$  of special  $\Delta$ -sets in  $\tilde{V}$  such that each  $\tilde{v} \in \tilde{V}$  is contained in at most  $\Delta$  special sets and a family  $\mathcal{B} \subseteq \binom{V}{\Delta}$  of forbidden  $\Delta$ -sets in  $V$  with  $|\mathcal{B}| \leq \mu|V|^\Delta$ . Then there is an embedding of  $H$  into  $G$  such that no special set is mapped to a forbidden set.*

At first sight, the rôle of the integer  $r$  in Lemma 12 (and also in Lemma 13 below) seems a little obscure. The only reason for stating the lemma as above is that it is more readily applicable in this form, since we will need it for pairs of partition classes  $(U, V)$  whose size in relation to  $n$  will be determined by the regularity lemma.

Our last main lemma, the connection lemma (Lemma 13), embeds graphs  $H$  into graphs  $G$  forming a system of  $p$ -dense pairs. In contrast to the blow-up lemma, however, the graph  $H$  has to be much smaller than the graph  $G$  now (see condition (A)). In addition, each vertex  $\tilde{y}$  of  $H$  is equipped with a candidate set  $C(\tilde{y})$  in  $G$  from which the connection lemma will choose the image of  $\tilde{y}$  in the embedding. Lemma 13 requires that these candidate sets are big (condition (D)) and that pairs of candidate sets that correspond to an edge of  $H$  form  $p$ -dense pairs (condition (E)). The remaining conditions ((B) and (C)) are conditions on the neighbourhoods and degrees of the vertices in  $H$  (with respect to the given partition of  $H$ ). For their statement we need the following additional definition.

For a graph  $H$  on vertex set  $\tilde{V} = \tilde{V}_1 \dot{\cup} \dots \dot{\cup} \tilde{V}_t$  and  $y \in \tilde{V}_i$  with  $i \in [t]$  define the *left degree* of  $y$  with respect to the partition  $\tilde{V}_1 \dot{\cup} \dots \dot{\cup} \tilde{V}_t$  to be  $\text{ldeg}(y; \tilde{V}_1, \dots, \tilde{V}_t) := \sum_{j=1}^{i-1} \deg_{\tilde{V}_j}(y)$ . When clear from the context we may also omit the partition and simply write  $\text{ldeg}(y)$ . For two sets of vertices  $S, T$  we denote the *joint neighbourhood* of (the vertices of)  $S$  in  $T$  by  $N_T^\cap(S) := \bigcap_{s \in S} N_T(s)$ .

**Lemma 13** (Connection lemma). *For all integers  $\Delta > 1$ ,  $t > 0$  and reals  $d > 0$  there are  $\varepsilon, \xi > 0$  such that for all positive integers  $r_1$  there is  $c > 1$  such that for all integers  $1 \leq r \leq r_1$  the following holds a.a.s. for  $\Gamma = \mathcal{G}_{n,p}$  with  $p \geq c(\log n/n)^{1/\Delta}$ . Let  $G \subseteq \Gamma$  be any graph on vertex set  $W = W_1 \dot{\cup} \dots \dot{\cup} W_t$  and let  $H$  be any graph on vertex set  $\tilde{W} = \tilde{W}_1 \dot{\cup} \dots \dot{\cup} \tilde{W}_t$ . Suppose further that for each  $i \in [t]$  each vertex  $\tilde{w} \in \tilde{W}_i$  is equipped with an arbitrary set  $X_{\tilde{w}} \subseteq V(\Gamma) \setminus W$  with the property that the indexed set system  $(X_{\tilde{w}}; \tilde{w} \in \tilde{W}_i)$  consists of pairwise disjoint sets such that the following holds. We define the *external degree* of  $\tilde{w}$  to be  $\text{edeg}(\tilde{w}) := |X_{\tilde{w}}|$ , its candidate set  $C(\tilde{w}) \subseteq W_i$  to be  $C(\tilde{w}) := N_{W_i}^\cap(X_{\tilde{w}})$ , and require that*

- (A)  $|W_i| \geq n/r$  and  $|\tilde{W}_i| \leq \xi n/r$ ,
- (B)  $\tilde{W}_i$  is a 3-independent set in  $H$ ,
- (C)  $\text{edeg}(\tilde{w}) + \text{ldeg}(\tilde{w}) = \text{edeg}(\tilde{v}) + \text{ldeg}(\tilde{v})$  and  $\deg_H(\tilde{w}) + \text{edeg}(\tilde{w}) \leq \Delta$  for all  $\tilde{w}, \tilde{v} \in \tilde{W}_i$ ,
- (D)  $|C(\tilde{w})| \geq ((d - \varepsilon)p)^{\text{edeg}(\tilde{w})} |W_i|$  for all  $\tilde{w} \in \tilde{W}_i$ , and
- (E)  $(C(\tilde{w}), C(\tilde{v}))$  forms an  $(\varepsilon, d, p)$ -dense pair for all  $\tilde{w}\tilde{v} \in E(H)$ .

*Then there is an embedding of  $H$  into  $G$  such that every vertex  $\tilde{w} \in \tilde{W}$  is mapped to a vertex in its candidate set  $C(\tilde{w})$ .*

The proof of this lemma is inherent in [23]. We adapt it to our setting in Section A.

## 6. STARS IN RANDOM GRAPHS

In this section we formulate two lemmas concerning properties of random graphs that will be useful when analysing neighbourhood properties of  $p$ -dense pairs in the following section. More precisely, we consider the following question here. Given a set of vertices  $X$  in a random graph  $\Gamma = \mathcal{G}_{n,p}$  together with a family  $\mathcal{F}$  of pairwise disjoint  $\ell$ -sets in  $V(\Gamma)$ . Then we would like to determine how many pairs  $(x, F)$  with  $x \in X$  and  $F \in \mathcal{F}$  have the property that  $x$  lies in the common neighbourhood of the vertices in  $F$ .

**Definition 14** (stars). *Let  $G = (V, E)$  be a graph,  $X$  be a subset of  $V$  and  $\mathcal{F}$  be a family of pairwise disjoint  $\ell$ -sets in  $V \setminus X$  for some  $\ell$ . Then the number of stars in  $G$  between  $X$  and  $\mathcal{F}$  is*

$$\# \text{stars}^G(X, \mathcal{F}) := \left| \{ (x, F) : x \in X, F \in \mathcal{F}, F \subseteq N_G(x) \} \right|. \quad (2)$$

Observe that in a random graph  $\Gamma = \mathcal{G}_{n,p}$  and for fixed sets  $X$  and  $\mathcal{F}$  the random variable  $\# \text{stars}^\Gamma(X, \mathcal{F})$  has binomial distribution  $\text{Bi}(|X||\mathcal{F}|, p^\ell)$ . This will be used in the proofs of the following lemmas. The first of these lemmas states that in  $\mathcal{G}_{n,p}$  the number of stars between  $X$  and  $\mathcal{F}$  does not exceed its expectation by more than seven times as long as  $X$  and  $\mathcal{F}$  are not too small. This is a straight-forward consequence of Chernoff's inequality.

**Lemma 15** (star lemma for big sets). *For all positive integers  $\Delta$ , and positive reals  $\nu$  there is  $c$  such that if  $p \geq c(\log n/n)^{1/\Delta}$  the following holds a.a.s. for  $\Gamma = \mathcal{G}_{n,p}$  on vertex set  $V$ . Let  $X$  be any subset of  $V$  and  $\mathcal{F}$  be any family of pairwise disjoint  $\Delta$ -sets in  $V \setminus X$ . If  $\nu n \leq |X| \leq |\mathcal{F}| \leq n$ , then*

$$\# \text{stars}^\Gamma(X, \mathcal{F}) \leq 7p^\Delta |X||\mathcal{F}|$$

*Proof.* Given  $\Delta$  and  $\nu$  let  $c$  be such that  $7c^\Delta \nu^2 \geq 3\Delta$ . From Chernoff's inequality (see [19, Chapter 2]) we know that  $\mathbb{P}[Y \geq 7\mathbb{E}Y] \leq \exp(-7\mathbb{E}Y)$  for a binomially distributed random variable  $Y$ . We conclude that for fixed  $X$  and  $\mathcal{F}$

$$\begin{aligned} \mathbb{P}[\# \text{stars}^\Gamma(X, \mathcal{F}) > 7p^\Delta |X||\mathcal{F}|] &\leq \exp(-7p^\Delta |X||\mathcal{F}|) \\ &\leq \exp(-7c^\Delta (\log n/n) \nu^2 n^2) \leq \exp(-3\Delta n \log n) \end{aligned}$$

by the choice of  $c$ . Thus the probability that there are sets  $X$  and  $\mathcal{F}$  violating the assertion of the lemma is at most

$$2^n n^{\Delta n} \exp(-3\Delta n \log n) \leq \exp(2\Delta n \log n - 3\Delta n \log n)$$

which tends to 0 as  $n$  tends to infinity.  $\square$

We will also need a variant of this lemma for smaller sets  $X$  and families  $\mathcal{F}$  which is provided in the next lemma. As a trade-off the bound on the number of stars provided by this lemma will be somewhat worse. This lemma almost appears in this form in [23]. The only (slight) modification that we need here is that  $X$  is allowed to be bigger than  $\mathcal{F}$ . However, the same proof as presented in [23] still works for this modified version. We delay it to Section B.1.

**Lemma 16** (star lemma for small sets). *For all positive integers  $\Delta$  and positive reals  $\xi$  there are positive constants  $\nu$  and  $c$  such that if  $p \geq c(\log n/n)^{1/\Delta}$ , then the following holds a.a.s. for  $\Gamma = \mathcal{G}_{n,p}$  on vertex set  $V$ . Let  $X$  be any subset of  $V$  and  $\mathcal{F}$  be any family of pairwise disjoint  $\Delta$ -sets in  $V \setminus X$ . If  $|X| \leq \nu n p^\Delta |\mathcal{F}|$  and  $|X|, |\mathcal{F}| \leq \xi n$ , then*

$$\# \text{stars}^\Gamma(X, \mathcal{F}) \leq p^\Delta |X||\mathcal{F}| + 6\xi n p^\Delta |\mathcal{F}|. \quad (3)$$

## 7. JOINT NEIGHBOURHOODS IN $p$ -DENSE PAIRS

As discussed in Section 4 it follows directly from the definition of  $p$ -denseness that sub-pairs of dense pairs form again dense pairs. In order to apply Lemma 12 and Lemma 13 together, we will need corresponding results on joint neighbourhoods in systems of dense pairs (see Lemmas 18 and 21). For this it is necessary to first introduce some notation.

Let  $G = (V, E)$  be a graph,  $\ell, T > 0$  be integers,  $p, \varepsilon, d$  be positive reals, and  $X, Y, Z \subseteq V$  be disjoint vertex sets. Recall that for a set  $B$  of vertices from  $V$  and a vertex set  $Y \subseteq V$  we call the set  $N_Y^\cap(B) = \bigcap_{b \in B} N_Y(b)$  the *joint neighbourhood* of (the vertices in)  $B$  in  $Y$ .

**Definition 17** (Bad and good vertex sets). *Let  $G, \ell, T, p, \varepsilon, d, X, Y$ , and  $Z$  be as above. We define the following family of  $\ell$ -sets in  $Y$  with small joint neighbourhood in  $Z$ :*

$$\text{bad}_{\varepsilon, d, p}^{G, \ell}(Y, Z) := \left\{ B \in \binom{Y}{\ell} : |N_Z^\cap(B)| < (d - \varepsilon)^\ell p^\ell |Z| \right\}. \quad (4)$$

If  $(Y, Z)$  has  $p$ -density  $d_{G,p}(Y, Z) \geq d - \varepsilon$ , then all  $\ell$ -sets  $T \in \binom{Y}{\ell}$  that are not in  $\text{bad}_{\varepsilon, d, p}^{G, \ell}(Y, Z)$  are called  $p$ -good in  $(Y, Z)$ . Let further

$$\text{Bad}_{\varepsilon, d, p}^{G, \ell}(X, Y, Z)$$

be the family of  $\ell$ -sets  $B \in \binom{X}{\ell}$  that contain an  $\ell'$ -set  $B' \subseteq B$  with  $\ell' > 0$  such that either  $|N_Y^\cap(B')| < (d - \varepsilon)^{\ell'} p^{\ell'} |Y|$  or  $(N_Y^\cap(B'), Z)$  is not  $(\varepsilon, d, p)$ -dense in  $G$ .

The following lemma states that  $p$ -dense pairs in random graphs have the property that most  $\ell$ -sets have big common neighbourhoods. Results of this type (with a slightly smaller exponent in the edge probability  $p$ ) were established in [21]. The proof of Lemma 18 is given in Section B.2.

**Lemma 18** (joint neighbourhood lemma). *For all integers  $\Delta, \ell \geq 1$  and positive reals  $d, \varepsilon'$  and  $\mu$ , there is  $\varepsilon > 0$  such that for all  $\xi > 0$  there is  $c > 1$  such that if  $p \geq c(\log n/n)^{1/\Delta}$ , then the following holds a.a.s. for  $\Gamma = \mathcal{G}_{n,p}$ . For  $n_1 \geq \xi p^{\Delta-1}n$ ,  $n_2 \geq \xi p^{\Delta-\ell}n$  let  $G = (X \dot{\cup} Y, E)$  be any bipartite subgraph of  $\Gamma$  with  $|X| = n_1$  and  $|Y| = n_2$ . If  $(X, Y)$  is an  $(\varepsilon, d, p)$ -dense pair, then  $|\text{bad}_{\varepsilon', d, p}^{G, \ell}(X, Y)| \leq \mu n_1^\ell$ .*

Thus we know that typical vertex sets in dense pairs inside random graphs are  $p$ -good. In the next lemma we observe that families of such  $p$ -good vertex sets exhibit strong expansion properties.

Given  $\ell$  and  $p$  we say that a bipartite graph  $G = (X \dot{\cup} Y, E)$  is  $(A, f)$ -expanding, if, for any family  $\mathcal{F} \subseteq \binom{X}{\ell}$  of pairwise disjoint  $p$ -good  $\ell$ -sets in  $(X, Y)$  with  $|\mathcal{F}| \leq A$ , we have  $|N_Y^\cap(\mathcal{F})| \geq f|\mathcal{F}|$ .

**Lemma 19** (expansion lemma). *For all positive integers  $\Delta$  and positive reals  $d$  and  $\varepsilon$ , there are positive  $\nu$  and  $c$  such that if  $p \geq c(\log n/n)^{1/\Delta}$ , then the following holds a.a.s. for  $\Gamma = \mathcal{G}_{n,p}$ . Let  $G = (X \dot{\cup} Y, E)$  be a bipartite subgraph of  $\Gamma$ . If  $(X, Y)$  is an  $(\varepsilon, d, p)$ -dense pair, then  $(X, Y)$  is  $(1/p^\Delta, \nu np^\Delta)$ -expanding.*

*Proof.* Given  $\ell, d, \varepsilon$ , set  $\delta := d - \varepsilon$ ,  $\xi := \delta^\Delta/7$  and let  $\nu'$  and  $c$  be the constants from Lemma 16 for this  $\Delta$  and  $\xi$ . Further, choose  $\nu$  such that  $\nu \leq \xi$  and  $\nu \leq \nu'$ . Let  $\mathcal{F} \subseteq \binom{X}{\Delta}$  be a family of pairwise disjoint  $p$ -good  $\Delta$ -sets with  $|\mathcal{F}| \leq 1/p^\Delta$ . Let  $U = N_Y^\cap(\mathcal{F})$  be the joint neighbourhood of  $\mathcal{F}$  in  $Y$ . We wish to show that  $|U| \geq (\nu np^\Delta)|\mathcal{F}|$ . Suppose the contrary. Then  $|U| < \nu' np^\Delta |\mathcal{F}|$ ,  $|U| < \nu np^\Delta |\mathcal{F}| \leq \nu n \leq \xi n$  and  $|\mathcal{F}| \leq 1/p^\Delta \leq c^\Delta n / \log n \leq \xi n$  for  $n$  sufficiently large and so we can apply Lemma 16 with parameters  $\Delta$  and  $\xi$  to  $U$  and  $\mathcal{F}$ . Since every member of  $\mathcal{F}$  is  $p$ -good in  $(X, Y)$ , we thus have

$$\begin{aligned} \delta^\Delta p^\Delta n |\mathcal{F}| &\leq \# \text{stars}^G(U, \mathcal{F}) \leq \# \text{stars}^\Gamma(U, \mathcal{F}) \stackrel{(3)}{\leq} p^\Delta |U| |\mathcal{F}| + 6\xi np^\Delta |\mathcal{F}| \\ &< p^\Delta (\nu np^\Delta) |\mathcal{F}| |\mathcal{F}| + 6\xi np^\Delta |\mathcal{F}| \leq \nu np^\Delta |\mathcal{F}| + 6\xi np^\Delta |\mathcal{F}| \leq 7\xi np^\Delta |\mathcal{F}|, \end{aligned}$$

which yields that  $\delta^\Delta < 7\xi$ , a contradiction.  $\square$

In the remainder of this section we are interested in the inheritance of  $p$ -denseness to sub-pairs  $(X', Y')$  of  $p$ -dense pairs  $(X, Y)$  in a graph  $G = (V, E)$ . It comes as a surprise that even for sets  $X'$  and  $Y'$  that are much smaller than the sets considered in the definition of  $p$ -denseness, such sub-pairs are typically dense. Phenomena of this type were observed in [21, 15].

Here, we will consider sub-pairs induced by neighbourhoods of vertices  $v \in V$  (which may or may not be in  $X \dot{\cup} Y$ ), i.e., sub-pairs  $(X', Y')$  where  $X'$  (or  $Y'$  or both) is the neighbourhood of  $v$  in  $Y$  (or in  $X$ ). Further, we only consider the case when  $G$  is a subgraph of a random graph  $\mathcal{G}_{n,p}$ .

In [23] an inheritance result of this form was obtained for triples of dense pairs. More precisely, the following holds for subgraphs  $G$  of  $\mathcal{G}_{n,p}$ . For sufficiently large vertex set  $X, Y$ , and  $Z$  in  $G$  such that  $(X, Y)$  and  $(Y, Z)$  form  $p$ -dense pairs we have that most vertices  $x \in X$  are such that  $(N_Y(x), Y)$  forms again a  $p$ -dense pair (with slightly changed parameters). If, moreover,  $(X, Z)$  forms a  $p$ -dense pair, too, then  $(N_Y(x), N_Z(x))$  is typically also a  $p$ -dense pair.

**Lemma 20** (inheritance lemma for vertices [23]). *For all integers  $\Delta > 0$  and positive reals  $d_0, \varepsilon'$  and  $\mu$  there is  $\varepsilon$  such that for all  $\xi > 0$  there is  $c > 1$  such that if  $p > c(\log n/n)^{1/\Delta}$ , then the following holds a.a.s. for  $\Gamma = \mathcal{G}_{n,p}$ . For  $n_1, n_3 \geq \xi p^{\Delta-1}n$  and  $n_2 \geq \xi p^{\Delta-2}n$  let  $G = (X \dot{\cup} Y \dot{\cup} Z, E)$  be any tripartite subgraph of  $\Gamma$  with  $|X| = n_1$ ,  $|Y| = n_2$ , and  $|Z| = n_3$ . If  $(X, Y)$  and  $(Y, Z)$*

are  $(\varepsilon, d, p)$ -dense pairs in  $G$  with  $d \geq d_0$ , then there are at most  $\mu n_1$  vertices  $x \in X$  such that  $(N(x) \cap Y, Z)$  is not an  $(\varepsilon', d, p)$ -dense pair in  $G$ .

If, additionally,  $(X, Z)$  is  $(\varepsilon, d, p)$ -dense and  $n_1, n_2, n_3 \geq \xi p^{\Delta-2} n$ , then there are at most  $\mu n_1$  vertices  $x \in X$  such that  $(N(x) \cap Y, N(x) \cap Z)$  is not an  $(\varepsilon', d, p)$ -dense pair in  $G$ .  $\square$

In order to combine the constrained blow-up lemma (Lemma 12) and the connection lemma (Lemma 13) in the proof of Theorem 1 we will need a version of this result for  $\ell$ -sets. Such a lemma, stating that joint neighbourhoods of certain  $\ell$ -sets form again  $p$ -dense pairs, can be obtained by an inductive argument from the first part of Lemma 20. We defer its proof to Section B.3.

**Lemma 21** (inheritance lemma for  $\ell$ -sets). *For all integers  $\Delta, \ell > 0$  and positive reals  $d_0, \varepsilon'$ , and  $\mu$  there is  $\varepsilon$  such that for all  $\xi > 0$  there is  $c > 1$  such that if  $p > c(\frac{\log n}{n})^{1/\Delta}$ , then the following holds a.a.s. for  $\Gamma = \mathcal{G}_{n,p}$ . For  $n_1, n_3 \geq \xi p^{\Delta-1} n$  and  $n_2 \geq \xi p^{\Delta-\ell-1} n$  let  $G = (X \dot{\cup} Y \dot{\cup} Z, E)$  be any tripartite subgraph of  $\Gamma$  with  $|X| = n_1$ ,  $|Y| = n_2$ , and  $|Z| = n_3$ . Assume further that  $(X, Y)$  and  $(Y, Z)$  are  $(\varepsilon, d, p)$ -dense pairs with  $d \geq d_0$ . Then*

$$|\text{Bad}_{\varepsilon', d, p}^{G, \ell}(X, Y, Z)| \leq \mu n_1^\ell.$$

## 8. PROOF OF THEOREM 1

In this section we present a proof of Theorem 1 that combines our four main lemmas, namely the lemma for  $G$  (Lemma 10), the lemma for  $H$  (Lemma 11), the constrained blow-up lemma (Lemma 12), and the connection lemma (Lemma 13). This proof follows the outline given in Section 5. In addition we will apply the inheritance lemma for  $\ell$ -sets (Lemma 21), which supplies an appropriate interface between the constrained blow-up lemma and the connection lemma.

*Proof of Theorem 1.* We first set up the constants. Given  $\eta, \gamma$ , and  $\Delta$  let  $t$  be the constant promised by the lemma for  $H$  (Lemma 11) for input  $\Delta$ . Set

$$\eta_G := \eta/10, \quad \text{and} \quad r_0 = 1, \quad (5)$$

and apply the lemma for  $G$  (Lemma 10) with input  $t, r_0, \eta_G$ , and  $\gamma$  in order to obtain  $\eta'_G$  and  $d$ . Next, the connection lemma (Lemma 13) with input  $\Delta, 2t$ , and  $d$  provides us with  $\varepsilon_{\text{CL}}$ , and  $\xi_{\text{CL}}$ . We apply the constrained blow-up lemma (Lemma 12) with  $\Delta, d$ , and  $\eta/2$  in order to obtain  $\varepsilon_{\text{BL}}$  and  $\mu_{\text{BL}}$ . With this we set

$$\eta_H := \min\{\eta/10, \xi_{\text{CL}}\eta'_G, 1/(\Delta+1)\}. \quad (6)$$

Choose  $\mu > 0$  such that

$$100t^2\mu \leq \eta_{\text{BL}}, \quad (7)$$

and apply Lemma 21 with  $\Delta$  and  $\ell = \Delta - 1$ ,  $d_0 = d$ ,  $\varepsilon' = \varepsilon_{\text{CL}}$ , and  $\mu$  to obtain  $\varepsilon_{21}$ . Let

$$\xi_{21} := \eta'_G/2r \quad (8)$$

and continue the application of Lemma 21 with  $\xi_{21}$  to obtain  $c_{21}$ . Now we can fix

$$\varepsilon := \min\{\varepsilon_{\text{CL}}, \varepsilon_{\text{BL}}, \varepsilon_{21}\} \quad (9)$$

and continue the application of Lemma 10 with input  $\varepsilon$  to get  $r_1$ . Let  $\hat{r}_{\text{BL}}$  and  $\hat{r}_{\text{CL}}$  be such that

$$\frac{2r_1}{1 - \eta_G} \leq \hat{r}_{\text{BL}} \quad \text{and} \quad \frac{2r_1}{\eta_G} \leq \hat{r}_{\text{CL}} \quad (10)$$

and let  $c_{\text{CL}}$  and  $c_{\text{BL}}$  be the constants obtained from the continued application of Lemma 13 with  $r_1$  replaced by  $\hat{r}_{\text{CL}}$  and Lemma 12 with  $r_1$  replaced by  $\hat{r}_{\text{BL}}$ , respectively.

We continue the application of Lemma 11 with input  $\eta_H$ . For each  $r \in [r_1]$  Lemma 11 provides a value  $\beta_r$ , among all of which we choose the smallest one and set  $\beta$  to this value. Finally, we set  $c := \max\{c_{\text{BL}}, c_{\text{CL}}, c_{21}\}$ .

Consider a graph  $\Gamma = \mathcal{G}_{n,p}$  with  $p \geq c(\log n/n)^{1/\Delta}$ . Then  $\Gamma$  a.a.s. satisfies the properties stated in Lemma 10, Lemma 12, Lemma 13, and Lemma 21, with the parameters previously specified. We assume in the following that this is the case and show that then also the following holds. For all subgraphs  $G \subseteq \Gamma$  and all graphs  $H$  such that  $G$  and  $H$  have the properties required by Theorem 1 we have  $H \subseteq G$ . To summarise the definition of the constants above, we can now assume that  $\Gamma$  satisfies the conclusion of the following lemmas:

- (L10) Lemma 10 for parameters  $t, r_0 = 1, \eta_G, \gamma, \eta'_G, d, \varepsilon$ , and  $r_1$ , i.e., if  $G$  is any spanning subgraph of  $\Gamma$  satisfying the requirements of Lemma 10, then we obtain a partition of  $G$  as specified in the lemma with these parameters,
- (L12) Lemma 12 for parameters  $\Delta, d, \eta/2, \varepsilon_{\text{BL}}, \mu_{\text{BL}}$ , and  $\hat{r}_{\text{BL}}$ ,
- (L13) Lemma 13 for parameters  $\Delta, 2t, d, \varepsilon_{\text{CL}}, \xi_{\text{CL}}$ , and  $\hat{r}_{\text{CL}}$ ,
- (L21) Lemma 21 for parameters  $\Delta, \ell = \Delta - 1, d_0 = d, \varepsilon' = \varepsilon_{\text{CL}}, \mu, \varepsilon_{21}$ , and  $\xi_{21}$ .

Now suppose we are given a graph  $G = (V, E) \subseteq \Gamma$  with  $\deg_G(v) \geq (\frac{1}{2} + \gamma) \deg_\Gamma(v)$  for all  $v \in V$  and  $|V| = n$ , and a graph  $H = (\tilde{V}, \tilde{E})$  with  $|\tilde{V}| = (1 - \eta)n$ . Before we show that  $H$  can be embedded into  $G$  we will use the lemma for  $G$  (Lemma 10) and the lemma for  $H$  (Lemma 11) to prepare  $G$  and  $H$  for this embedding.

First we use the fact that  $\Gamma$  has property (L10). Hence, for the graph  $G$  we obtain an  $r$  with  $1 \leq r \leq r_1$  from Lemma 10, together with a set  $V_0 \subseteq V$  with  $|V_0| \leq \varepsilon n$ , and a mapping  $g: V \setminus V_0 \rightarrow R_{r,t}$  such that (G1)–(G3) of Lemma 10 are fulfilled. For all  $i \in [r], j \in [2t]$  let  $U_i, V_i, C_{i,j}, C'_{i,j}, B_{i,j}$ , and  $B'_{i,j}$  be the sets defined in Lemma 10. Recall that these sets were called big clusters, connecting clusters, and balancing clusters. With this the graph  $G$  is prepared for the embedding. We now turn to the graph  $H$ .

We assume for simplicity that  $2r/(1 - \eta_G)$  and  $r/(\eta'_G)$  are integers and define

$$r_{\text{BL}} := 2r/(1 - \eta_G) \quad \text{and} \quad r_{\text{CL}} := 2r/\eta'_G. \quad (11)$$

We apply Lemma 11 which we already provided with  $\Delta$  and  $\eta_H$ . For input  $H$  this lemma provides a homomorphism  $h$  from  $H$  to  $R_{r,t}$  such that (H1)–(H5) of Lemma 11 are fulfilled. For all  $i \in [r], j \in [2t]$  let  $\tilde{U}_i, \tilde{V}_i, \tilde{C}_{i,j}, \tilde{C}'_{i,j}, \tilde{B}_{i,j}, \tilde{B}'_{i,j}$ , and  $\tilde{X}_i$  be the sets asserted by Lemma 11. Further, set  $C_i := C_{i,1} \dot{\cup} \dots \dot{\cup} C_{i,2t}$ ,  $\tilde{C}_i := \tilde{C}_{i,1} \dot{\cup} \dots \dot{\cup} \tilde{C}_{i,2t}$ , that is,  $C_i$  consists of connecting clusters and  $\tilde{C}_i$  of connecting vertices. Define  $C'_i, \tilde{C}'_i, B_i, \tilde{B}_i, B'_i$ , and  $\tilde{B}'_i$  analogously ( $B_i$  consists of balancing clusters and  $\tilde{B}_i$  of balancing vertices).

Our next goal will be to appeal to property (L12) which asserts that we can apply the constrained blow-up lemma (Lemma 12) for each  $p$ -dense pair  $(U_i, V_i)$  with  $i \in [r]$  individually and embed  $H[\tilde{U}_i \dot{\cup} \tilde{V}_i]$  into this pair. For this we fix  $i \in [r]$ . We will first set up special  $\Delta$ -sets  $\mathcal{H}_i$  and forbidden  $\Delta$ -sets  $\mathcal{B}_i$  for the application of Lemma 12. The idea is as follows. With the help of Lemma 12 we will embed all vertices in  $\tilde{U}_i \dot{\cup} \tilde{V}_i$ . But all connecting and balancing vertices of  $H$  remain unembedded. They will be handled by the connection lemma, Lemma 13, later on. However, these two lemmas cannot operate independently. If, for example, a connecting vertex  $\tilde{y}$  has three neighbours in  $\tilde{V}_i$ , then these neighbours will be already mapped to vertices  $v_1, v_2, v_3$  in  $V_i$  (by the blow-up lemma) when we want to embed  $\tilde{y}$ . Accordingly the image of  $\tilde{y}$  in the embedding is confined to the joint neighbourhood of the vertices  $v_1, v_2, v_3$  in  $G$ . In other words, this joint neighbourhood will be the candidate set  $C(\tilde{y})$  in the application of Lemma 13. This lemma requires, however, that candidate sets are not too small (condition (D) of Lemma 13) and, in addition, that candidate sets of any two adjacent vertices induce  $p$ -dense pairs (condition (E)). Hence we need to be prepared for these requirements. This will be done via the special and forbidden sets. The family of special sets  $\mathcal{H}_i$  will contain neighbourhoods in  $\tilde{V}_i$  of connecting or balancing vertices  $\tilde{y}$  of  $H$  (observe that such vertices do not have neighbours in  $\tilde{U}_i$ , see Figure 1). The family of forbidden sets  $\mathcal{B}_i$  will consist of sets in  $V_i$  which are “bad” for the embedding of these neighbourhoods in view of (D) and (E) of Lemma 13 (recall that Lemma 12 does not map special sets to forbidden sets). Accordingly,  $\mathcal{B}_i$  contains  $\Delta$ -sets that have small common neighbourhoods or do not induce  $p$ -dense pairs in one of the relevant balancing or connecting clusters. We will next give the details of this construction of  $\mathcal{H}_i$  and  $\mathcal{B}_i$ .

We start with the special  $\Delta$ -sets  $\mathcal{H}_i$ . As explained, we would like to include in the family  $\mathcal{H}_i$  all neighbourhoods of vertices  $\tilde{w}$  of vertices outside  $\tilde{U}_i \dot{\cup} \tilde{V}_i$ . Such neighbourhoods clearly lie entirely in the set  $\tilde{X}_i$  provided by Lemma 11. However, they need not necessarily be  $\Delta$ -sets (in fact, by (H4) of Lemma 11, they are of size at most  $\Delta - 1$ ). Therefore we have to “pad” these neighbourhoods in order to obtain  $\Delta$ -sets. This is done as follows. We start by picking an arbitrary set of  $\Delta|\tilde{X}_i|$  vertices (which will be used for the “padding”) in  $\tilde{V}_i \setminus \tilde{X}_i$ . We add these vertices to  $\tilde{X}_i$



and call the resulting set  $\tilde{X}'_i$ . This is possible because (H5) of Lemma 11 and (6) imply that  $|\tilde{X}'_i| \leq (\Delta + 1)|\tilde{X}_i| \leq (\Delta + 1)\eta_H|\tilde{V}_i| \leq |\tilde{V}_i|$ .

Now let  $\tilde{Y}_i$  be the set of vertices in  $\tilde{B}_i \dot{\cup} \tilde{C}_i$  with neighbours in  $\tilde{V}_i$ . These are the vertices for whose neighbourhoods we will include  $\Delta$ -sets in  $\mathcal{H}_i$ . It follows from the definition of  $\tilde{X}_i$  that  $|\tilde{Y}_i| \leq \Delta|\tilde{X}_i|$ . Let  $\tilde{y} \in \tilde{Y}_i \subseteq \tilde{B}_i \cup \tilde{C}_i$ . By the definition of  $\tilde{X}_i$  we have  $N_H(\tilde{y}) \subseteq \tilde{X}_i$ . Next, we let

$$\tilde{X}_{\tilde{y}} \text{ be the set of neighbours of } \tilde{y} \text{ in } \tilde{V}_i \quad (12)$$

As explained,  $\tilde{y}$  has strictly less than  $\Delta$  neighbours in  $\tilde{V}_i$  and hence we choose additional vertices from  $\tilde{X}'_i \setminus \tilde{X}_i$ . In this way we obtain for each  $\tilde{y} \in \tilde{Y}_i$  a  $\Delta$ -set  $N_{\tilde{y}} \in \tilde{X}'_i$  with

$$N_{\tilde{X}_i}(\tilde{y}) = N_{\tilde{V}_i}(\tilde{y}) = \tilde{X}_{\tilde{y}} \subseteq N_{\tilde{y}}. \quad (13)$$

We make sure, in this process, that for any two different  $\tilde{y}$  and  $\tilde{y}'$  we never include the same additional vertex from  $\tilde{X}'_i \setminus \tilde{X}_i$ . This is possible because  $|\tilde{X}'_i \setminus \tilde{X}_i| \geq \Delta|\tilde{X}_i| \geq |\tilde{Y}_i|$ . We can thus guarantee that

$$\text{each vertex in } \tilde{X}'_i \text{ is contained in at most } \Delta \text{ sets } N_{\tilde{y}}. \quad (14)$$

The family of special  $\Delta$ -sets for the application of Lemma 12 on  $(U_i, V_i)$  is then

$$\mathcal{H}_i := \{N_{\tilde{y}} : \tilde{y} \in \tilde{Y}_i\}. \quad (15)$$

Note that this is indeed a family of  $\Delta$ -sets encoding all neighbourhoods in  $\tilde{U}_i \dot{\cup} \tilde{V}_i$  of vertices outside this set.

Now we turn to the family  $\mathcal{B}_i$  of forbidden  $\Delta$ -sets. Recall that this family should contain sets that are forbidden for the embedding of the special  $\Delta$ -sets because their joint neighbourhood in a (relevant) balancing or connecting cluster is small or does not induce a  $p$ -dense pair. More precisely, we are interested in  $\Delta$ -sets  $S$  that have one of the following properties. Either  $S$  has a small common neighbourhood in some cluster from  $B_i$  or from  $C_i$  (observe that only balancing vertices from  $\tilde{B}_i$  and connecting vertices from  $\tilde{C}_i$  have neighbours in  $\tilde{V}_i$ ). Or the neighbourhood  $N_D^\cap(S)$  of  $S$  in a cluster  $D$  from  $B_i$  or  $C_i$ , respectively, is such that  $(N_D^\cap(S), D')$  is not  $p$ -dense for some cluster  $D'$  from  $B'_i \cup B'_{i+1}$  or  $C'_i \cup C'_{i+1}$  (observe that edges between balancing vertices run only between  $\tilde{B}_i$  and  $\tilde{B}'_i \cup \tilde{B}'_{i+1}$  and edges between connecting vertices only between  $\tilde{C}_i$  and  $\tilde{C}'_i \cup \tilde{C}'_{i+1}$ ).

For technical reasons, however, we need to digress from this strategy slightly: We want to bound the number of  $\Delta$ -sets in  $\mathcal{B}_i$  with the help of the inheritance lemma for  $\ell$ -sets, Lemma 21, later. Notice that, thanks to the lower bound on  $n_2$  in Lemma 21, this lemma cannot be applied (in a meaningful way) for  $\Delta$ -sets. But it can be applied for  $(\Delta - 1)$ -sets. Therefore, we will not consider  $\Delta$ -sets directly but first construct an auxiliary family of  $(\Delta - 1)$ -sets and then, again, “pad” these sets to obtain a family of  $\Delta$ -sets. Observe that the strategy outlined while setting up the special sets  $\mathcal{H}_i$  still works with these  $(\Delta - 1)$ -sets: neighbourhoods of connecting or balancing vertices in  $\tilde{V}_i$  are of size at most  $\Delta - 1$  by (H4) of Lemma 11.

But now let us finally give the details. We first define the auxiliary family of  $(\Delta - 1)$ -sets as follows:

$$\begin{aligned} \mathcal{B}'_i := & \bigcup_{\substack{i' \in \{i, i+1\}, j, j' \in [2t] \\ (c_{i,j}, c'_{i',j'}) \in R_{r,t}}} \text{Bad}_{\varepsilon_{\text{cl}}, d, p}^{G, \Delta-1}(V_i, C_{i,j}, C'_{i',j'}) \cup \\ & \bigcup_{\substack{j, j' \in [2t] \\ (b_{i,j}, b'_{i,j'}) \in R_{r,t}}} \text{Bad}_{\varepsilon_{\text{cl}}, d, p}^{G, \Delta-1}(V_i, B_{i,j}, B'_{i,j'}). \end{aligned} \quad (16)$$

We will next bound the size of this family by appealing to property (L21), and hence Lemma 21, with the tripartite graphs  $G[V_i, C_{i,j}, C'_{i',j'}]$  and  $G[V_i, B_{i,j}, B'_{i,j'}]$  with indices as in the definition of  $\mathcal{B}'_i$ . For this we need to check the conditions appearing in this lemma. By the definition of  $R_{r,t}$  and (G3) of Lemma 10 all pairs  $(C_{i,j}, C'_{i',j'})$  and  $(B_{i,j}, B'_{i,j'})$  appearing in the definition of  $\mathcal{B}'_i$  as well as the pairs  $(V_i, C_{i,j})$  and  $(V_i, B_{i,j})$  with  $j \in [2t]$  are  $(\varepsilon, d, p)$ -dense. For the vertex sets of these



dense pairs we know  $|V_i|, |C'_{i',j'}|, |B'_{i,j'}| \geq \eta'_G n/2r \geq \xi_{21} p^{\Delta-1} n$  and  $|C_{i,j}|, |B_{i,j}| \geq \eta'_G n/2r = \xi_{21} n$  by (G1) and (G2) of Lemma 10 and (8). Thus, since  $\varepsilon \leq \varepsilon_{21}$ , property (L21) implies that the family

$$\text{Bad}_{\varepsilon_{\text{CL}}, d, p}^{G, \Delta-1}(V_i, C_{i,j}, C'_{i',j'}), \text{ and } \text{Bad}_{\varepsilon_{\text{CL}}, d, p}^{G, \Delta-1}(V_i, B_{i,j}, B'_{i,j'})$$

is of size  $\mu|V_i|^{\Delta-1}$  at most. It follows from (16) that  $|\mathcal{B}'_i| \leq 8t^2 \mu|V_i|^{\Delta-1}$  which is at most  $\mu_{\text{BL}}|V_i|^{\Delta-1}$  by (7). The family of forbidden  $\Delta$ -sets is then defined by

$$\mathcal{B}_i := \mathcal{B}'_i \times V_i \quad \text{and we have} \quad |\mathcal{B}_i| \leq \mu_{\text{BL}}|V_i|^\Delta. \quad (17)$$

Having defined the special and forbidden  $\Delta$ -sets we are now ready to appeal to (L12) and use the constrained blow-up lemma (Lemma 12) with parameters  $\Delta, d, \eta/2, \varepsilon_{\text{BL}}, \mu_{\text{BL}}, \hat{r}_{\text{BL}}$ , and  $r_{\text{BL}}$  separately for each pair of graphs  $G_i := (U_i, V_i)$  and  $H_i := H[\widetilde{U}_i \dot{\cup} \widetilde{V}_i]$ . Let us quickly check that the constant  $r_{\text{BL}}$  and the graphs  $G_i$  and  $H_i$  satisfy the required conditions. Observe first, that  $1 \leq r_{\text{BL}} = 2r/(1 - \eta_G) \leq 2r_1/(1 - \eta_G) \leq \hat{r}_{\text{BL}}$  by (11) and (10). Moreover  $(U_i, V_i)$  is an  $(\varepsilon_{\text{BL}}, d, p)$ -dense pair by (G3) of Lemma 10 and (9). (G1) implies

$$|U_i| \geq (1 - \eta_G) \frac{n}{2r} \stackrel{(11)}{=} \frac{n}{r_{\text{BL}}}$$

and similarly  $|V_i| \geq n/r_{\text{BL}}$ . By (H1) of Lemma 11 we have

$$\begin{aligned} |\widetilde{U}_i| &\leq (1 + \eta_H) \frac{n}{2r} \leq (1 + \eta_H)(1 - \eta) \frac{n}{2r} \leq (1 + \eta_H - \eta) \frac{n}{2r} \stackrel{(5),(6)}{\leq} (1 - \tfrac{1}{2}\eta - \eta_G) \frac{n}{2r} \\ &\leq (1 - \tfrac{1}{2}\eta)(1 - \eta_G) \frac{n}{2r} \stackrel{(11)}{=} (1 - \tfrac{1}{2}\eta) \frac{n}{r_{\text{BL}}} \end{aligned}$$

and similarly  $|\widetilde{V}_i| \leq (1 - \tfrac{\eta}{2})n/r_{\text{BL}}$ . For the application of Lemma 12, let the families of special and forbidden  $\Delta$ -sets be defined in (15) and (17), respectively. Observe that (14) and (17) guarantee that the required conditions (of Lemma 12) are satisfied. Consequently there is an embedding of  $H_i$  into  $G_i$  for each  $i \in [r]$  such that no special  $\Delta$ -set is mapped to a forbidden  $\Delta$ -set. Denote the united embedding resulting from these  $r$  applications of the constrained blow-up lemma by  $f_{\text{BL}} : \bigcup_{i \in [r]} \widetilde{U}_i \cup \widetilde{V}_i \rightarrow \bigcup_{i \in [r]} U_i \cup V_i$ .

It remains to verify that  $f_{\text{BL}}$  can be extended to an embedding of all vertices of  $H$  into  $G$ . We still need to take care of the balancing and connecting vertices. For this purpose we will, again, fix  $i \in [r]$  and use property (L13) which states that the conclusion of the connection lemma (Lemma 13) holds for parameters  $\Delta, 2t, d, \varepsilon_{\text{CL}}, \xi_{\text{CL}}$ , and  $\hat{r}_{\text{CL}}$ . We will apply this lemma with input  $r_{\text{CL}}$  to the graphs  $G'_i := G[W_i]$  and  $H'_i := H[\widetilde{W}_i]$  where  $W_i$  and  $\widetilde{W}_i$  and their partitions for the application of the connection lemma are as follows (see Figure 2). Let  $W_i := W_{i,1} \dot{\cup} \dots \dot{\cup} W_{i,8t}$  where for all  $j \in [t], k \in [2t]$  we set

$$\begin{aligned} W_{i,j} &:= C_{i,t+j}, & W_{i,2t+j} &:= C_{i+1,j}, & W_{i,2t+j} &:= C'_{i,t+j}, \\ W_{i,3t+j} &:= C'_{i+1,j}, & W_{i,4t+k} &:= B_{i,k}, & W_{i,6t+k} &:= B'_{i,k}. \end{aligned}$$

(This means that we propose the clusters in the following order to the connection lemma. The connecting clusters without primes come first, then the connecting clusters with primes, then the balancing clusters without primes, and finally the balancing clusters with primes.)

The partition  $\widetilde{W}_i := \widetilde{W}_{i,1} \dot{\cup} \dots \dot{\cup} \widetilde{W}_{i,8t}$  of the vertex set  $\widetilde{W}_i$  of  $H'_i$  is defined accordingly, i.e., for all  $j \in [t], k \in [2t]$  we set

$$\begin{aligned} \widetilde{W}_{i,j} &:= \widetilde{C}_{i,t+j}, & \widetilde{W}_{i,2t+j} &:= \widetilde{C}_{i+1,j}, & \widetilde{W}_{i,2t+j} &:= \widetilde{C}'_{i,t+j}, \\ \widetilde{W}_{i,3t+j} &:= \widetilde{C}'_{i+1,j}, & \widetilde{W}_{i,4t+k} &:= \widetilde{B}_{i,k}, & \widetilde{W}_{i,6t+k} &:= \widetilde{B}'_{i,k}. \end{aligned}$$

To check whether we can apply the connecting lemma observe first that

$$1 \leq 2r/\eta_G \leq 2r_1/\eta_G \leq \hat{r}_{\text{CL}}$$




FIGURE 2. The partition  $W_i = W_{i,1} \dot{\cup} \dots \dot{\cup} W_{i,8t}$  of  $G'_i = G[W_i]$  for the special case  $t = 2$ .

by (10). For  $\tilde{y} \in \tilde{W}_{i,j}$  with  $j \in [8t]$  recall from (12) (using that each vertex in  $H$  has neighbours in at most one set  $\tilde{V}_{i'}$ , see Figure 1) that

$$\tilde{X}_{\tilde{y}} \text{ is the set of neighbours of } \tilde{y} \text{ in } \tilde{V}_i \cup \tilde{V}_{i+1} \text{ and set } X_{\tilde{y}} := f_{\text{BL}}(\tilde{X}_{\tilde{y}}). \quad (18)$$

Then the indexed set system  $(\tilde{X}_{\tilde{y}} : \tilde{y} \in \tilde{W}_{i,j})$  consists of pairwise disjoint sets because  $\tilde{W}_{i,j}$  is 3-independent in  $H$  by (H3) of Lemma 11. Thus also  $(X_{\tilde{y}} : \tilde{y} \in \tilde{W}_{i,j})$  consists of pairwise disjoint sets, as required by Lemma 13. Now let the external degree and the candidate set of  $\tilde{y} \in \tilde{W}_{i,j}$  be defined as in Lemma 13, i.e.,

$$\text{edeg}(\tilde{y}) := |X_{\tilde{y}}| \quad \text{and} \quad C(\tilde{y}) := N_{\tilde{W}_{i,j}}^\cap(X_{\tilde{y}}). \quad (19)$$

Observe that this implies  $C(\tilde{y}) = W_{i,j}$  if  $\tilde{X}_{\tilde{y}} = \emptyset$  and hence  $X_{\tilde{y}} = \emptyset$ . Now we will check that conditions (A)–(E) of Lemma 13 are satisfied. From (G2) of Lemma 10 and (H2) of Lemma 11 it follows that

$$\begin{aligned} |W_{i,j}| &\stackrel{\text{(G2)}}{\geq} \eta'_G \frac{n}{2r} \stackrel{\text{(11)}}{=} \frac{n}{r_{\text{CL}}} \quad \text{and} \\ |\tilde{W}_{i,j}| &\stackrel{\text{(H2)}}{\leq} \eta_H \frac{m}{2r} \leq \eta_H \frac{n}{2r} \stackrel{\text{(11)}}{=} \frac{\eta_H}{\eta'_G} \frac{n}{r_{\text{CL}}} \stackrel{\text{(6)}}{\leq} \xi_{\text{CL}} \frac{n}{r_{\text{CL}}} \end{aligned}$$

and thus we have condition (A). By (H3) of Lemma 11 we also get condition (B) of Lemma 13. Further, it follows from (H4) of Lemma 11 that  $\text{edeg}(\tilde{y}) = \text{edeg}(\tilde{y}')$  and  $\text{ldeg}(\tilde{y}) = \text{ldeg}(\tilde{y}')$  for all  $\tilde{y}, \tilde{y}' \in \tilde{W}_{i,j}$  with  $j \in [8t]$ . In addition  $\Delta(H) \leq \Delta$  and hence

$$\begin{aligned} \deg_{H'_i}(\tilde{y}) + \text{edeg}(\tilde{y}) &\stackrel{\text{(19)}}{=} |N_{\tilde{W}_i}(\tilde{y})| + |X_{\tilde{y}}| \\ &\stackrel{\text{(18)}}{=} |N_{\tilde{W}_i}(\tilde{y})| + |N_{\tilde{V}_i \cup \tilde{V}_{i+1}}(\tilde{y})| \leq \deg_H(\tilde{y}) \leq \Delta \end{aligned}$$

and thus condition (C) of Lemma 13 is satisfied. To check conditions (D) and (E) of Lemma 13 observe that for all  $\tilde{y} \in \tilde{C}'_{i',j}$  with  $i' \in \{i, i+1\}$  and  $j \in [2t]$  we have  $C(\tilde{y}) = C'_{i',j}$  as  $\tilde{y}$  has no neighbours in  $\tilde{V}_i$  or  $\tilde{V}_{i+1}$  and hence the external  $\text{edeg}(\tilde{y}) = 0$  (see (18) and (19)). Thus (D) is satisfied for  $\tilde{y} \in \tilde{C}'_{i',j}$ , and similarly for  $\tilde{y} \in \tilde{B}'_{i',j}$ . For all  $\tilde{y} \in \tilde{C}_{i,j}$  with  $t < j \leq 2t$  on the other hand we have  $\tilde{X}_{\tilde{y}} \subseteq N_{\tilde{y}} \in \binom{\tilde{V}_i}{\Delta}$  by (12). Recall that  $N_{\tilde{y}}$  was a special  $\Delta$ -set in the application of the restricted blow-up lemma on  $G_i = (U_i, V_i)$  and  $H_i = H[\tilde{U}_i \dot{\cup} \tilde{V}_i]$  owing to (15). Therefore

$N_{\tilde{y}}$  is not mapped to a forbidden  $\Delta$ -set in  $\mathcal{B}_i \subseteq \binom{V_i}{\Delta}$  by  $f_{\text{BL}}$  and thus, by (16), to no  $\Delta$ -set in  $\text{Bad}_{\varepsilon_{\text{CL}}, d, p}^{G, \Delta-1}(V_i, C_{i,j}, C'_{i',j'}) \times V_i$  with  $i' \in \{i, i+1\}, j, j' \in [2t]$  and  $(c_{i,j}, c'_{i',j'}) \in R_{r,t}$ . We infer that the set  $f_{\text{BL}}(\tilde{X}_{\tilde{y}}) = X_{\tilde{y}} \in \binom{V_{i(\tilde{y})}}{\text{edeg}(\tilde{y})}$  satisfies  $|N_{C_{i,j}}^\cap(X_{\tilde{y}})| \geq (d - \varepsilon_{\text{CL}})^{\text{edeg}(\tilde{y})} p^{\text{edeg}(\tilde{y})} |C_{i,j}|$  and is such that

$$(N_{C_{i,j}}^\cap(X_{\tilde{y}}), C'_{i',j'}) \text{ is } (\varepsilon_{\text{CL}}, d, p)\text{-dense} \\ \text{for all } i' \in \{i, i+1\}, j, j' \in [2t] \text{ with } (c_{i,j}, c'_{i',j'}) \in R_{r,t}. \quad (20)$$

Since we chose  $C(\tilde{y}) = N^\cap(X_{\tilde{y}}) \cap C_{i,j}$  in (19) we get condition (D) of Lemma 13 also for  $\tilde{y} \in \tilde{C}_{i,j}$  with  $t < j \leq 2t$ . For  $\tilde{y} \in \tilde{C}_{i+1,j}$  with  $j \in [t]$  the same argument applies with  $\tilde{X}_{\tilde{y}} \subseteq N_{\tilde{y}} \in \binom{V_{i+1}}{\Delta^{i+1}}$ , and for  $\tilde{y} \in \tilde{B}_{i,j}$  with  $j \in [2t]$  the same argument applies with  $\tilde{X}_{\tilde{y}} \subseteq N_{\tilde{y}} \in \binom{V_i}{\Delta}$ .

Now it will be easy to see that we get (E) of Lemma 13. Indeed, recall again that  $C(\tilde{y}) = C'_{i',j'}$  for all  $\tilde{y} \in \tilde{C}'_{i',j'}$  and  $C(\tilde{y}) = B'_{i',j'}$  for all  $\tilde{y} \in \tilde{B}'_{i',j'}$  with  $i' \in \{i, i+1\}$  and  $j \in [2t]$ . In addition, the mapping  $h$  constructed by Lemma 11 is a homomorphism from  $H$  to  $R_{r,t}$ . Hence (20) and property (G3) of Lemma 10 assert that condition (E) of Lemma 13 is satisfied for all edges  $\tilde{y}\tilde{y}'$  of  $H'_i = H[\tilde{W}_i]$  with at least one end, say  $\tilde{y}$ , in a cluster  $\tilde{C}'_{i',j'}$  or  $\tilde{B}'_{i',j'}$ . This is true because then  $C(\tilde{y}) = W_{i,k}$  where  $\tilde{W}_{i,k}$  is the cluster containing  $\tilde{y}$ , and  $C(\tilde{y}') = N^\cap(X_{\tilde{y}'}) \cap W_{i,k'}$  where  $\tilde{W}_{i,k'}$  is the cluster containing  $\tilde{y}'$ . Moreover, since  $h$  is a homomorphism all edges  $\tilde{y}\tilde{y}'$  in  $H'_i = H[\tilde{W}_i]$  have at least one end in a cluster  $\tilde{C}'_{i',j'}$  or  $\tilde{B}'_{i',j'}$ .

So conditions (A)–(E) are satisfied and we can apply Lemma 13 to get embeddings of  $H'_i = H[\tilde{W}_i]$  into  $G'_i = G[W_i]$  for all  $i \in [r]$  that map vertices  $\tilde{y} \in \tilde{W}_i$  (i.e. connecting and balancing vertices) to vertices  $y \in W_i$  in their candidate sets  $C(\tilde{y})$ . Let  $f_{\text{CL}}$  be the united embedding resulting from these  $r$  applications of the connection lemma and denote the embedding that unites  $f_{\text{BL}}$  and  $f_{\text{CL}}$  by  $f$ .

To finish the proof we verify that  $f$  is an embedding of  $H$  into  $G$ . Let  $\tilde{x}\tilde{y}$  be an edge of  $H$ . By definition of the spin graph  $R_{r,t}$  and since the mapping  $h$  constructed by Lemma 11 is a homomorphism from  $H$  to  $R_{r,t}$  we only need to distinguish the following cases for  $i \in [r]$  and  $j, j' \in [2t]$  (see also Figure 1):

- case 1: If  $\tilde{x} \in \tilde{V}_i$  and  $\tilde{y} \in \tilde{U}_i$ , then  $f(\tilde{x}) = f_{\text{BL}}(\tilde{x})$  and  $f(\tilde{y}) = f_{\text{BL}}(\tilde{y})$  and thus the constrained blow-up lemma guarantees that  $f(\tilde{x})f(\tilde{y})$  is an edge of  $G_i$ .
- case 2: If  $\tilde{x} \in \tilde{W}_i$  and  $\tilde{y} \in \tilde{W}_i$ , then  $f(\tilde{x}) = f_{\text{CL}}(\tilde{x})$  and  $f(\tilde{y}) = f_{\text{CL}}(\tilde{y})$  and thus the connection lemma guarantees that  $f(\tilde{x})f(\tilde{y})$  is an edge of  $G'_i$ .
- case 3: If  $\tilde{x} \in \tilde{V}_i$  and  $\tilde{y} \in \tilde{W}_i$ , then either  $\tilde{y} \in \tilde{C}_{i,j}$  or  $\tilde{y} \in \tilde{B}_{i,j}$  for some  $j$ . Moreover,  $f(\tilde{x}) = f_{\text{BL}}(\tilde{x})$  and therefore by (19) the candidate set  $C(\tilde{y})$  of  $\tilde{y}$  satisfies  $C(\tilde{y}) \subseteq N_{C_{i,j}}(f(\tilde{x}))$  or  $C(\tilde{y}) \subseteq N_{B_{i,j}}(f(\tilde{x}))$ , respectively. As  $f(\tilde{y}) = f_{\text{CL}}(\tilde{y}) \in C(\tilde{y})$  we also get that  $f(\tilde{x})f(\tilde{y})$  is an edge of  $G$  in this case.

It follows that  $f$  maps all edges of  $H$  to edges of  $G$ , which finishes the proof of the theorem.  $\square$

## 9. A $p$ -DENSE PARTITION OF $G$

For the proof of the Lemma for  $G$  we shall apply the minimum degree version of the sparse regularity lemma (Lemma 8). Observe that this lemma guarantees that the reduced graph of the regular partition we obtain is dense. Thus we can apply Theorem 2 to this reduced graph. In the proof of Lemma 10 we use this theorem to find a copy of the ladder  $R_r^*$  in the reduced graph (the graphs  $R_r^*$  and  $R_{r,t}$  are defined in Section 5 on page 6, see also Figure 1). Then we further partition the clusters in this ladder to obtain a regular partition whose reduced graph contains a spin graph  $R_{r,t}$ . Recall that this partition will consist of a series of so-called *big clusters* which we denote by  $U_i$  and  $V_i$ , and a series of smaller clusters called *balancing clusters*  $B_{i,j}$ ,  $B'_{i,j}$  and *connecting clusters*  $C_{i,j}$ ,  $C'_{i,j}$  with  $i \in [r]$ ,  $j \in [2t]$ . We will now give the details.

*Proof of Lemma 10.* Given  $t, r_0, \eta$ , and  $\gamma$  choose  $\eta'$  such that

$$\frac{\eta}{5} + \left(\frac{4}{\gamma} + 2\right)t \cdot \eta' \leq \frac{\eta}{2} \quad (21)$$

and set  $d := \gamma/4$ . Apply Theorem 2 with input  $r_{\text{BK}} := 2, \Delta = 3$  and  $\gamma/2$  to obtain the constants  $\beta$  and  $k_{\text{BK}} := n_0$ . For input  $\varepsilon$  set

$$r'_0 := \max\{2r_0 + 1, k_{\text{BK}}, 3/\beta, 6/\gamma, 2/\varepsilon, 10/\eta\} \quad (22)$$

and choose  $\varepsilon'$  such that

$$\varepsilon'/\eta' \leq \varepsilon/2, \quad \text{and} \quad \varepsilon' \leq \min\{\gamma/4, \eta/10\}. \quad (23)$$

Lemma 8 applied with  $\alpha := \frac{1}{2} + \gamma, \varepsilon', r'_0$  then gives us the missing constant  $r_1$ .

Assume that  $\Gamma$  is a typical graph from  $\mathcal{G}_{n,p}$  with  $\log^4 n/(pn) = o(1)$ , in the sense that it satisfies the conclusion of Lemma 8, and let  $G = (V, E) \subseteq \Gamma$  satisfy  $\deg_G(v) \geq (\frac{1}{2} + \gamma)\deg_\Gamma(v)$  for all  $v \in V$ . Lemma 8 applied with  $\alpha = \frac{1}{2} + \gamma, \varepsilon', r'_0$ , and  $d$  to  $G$  gives us an  $(\varepsilon', d, p)$ -dense partition  $V = V'_0 \dot{\cup} V'_1 \dot{\cup} \dots \dot{\cup} V'_{r'}$  of  $G$  with reduced graph  $R'$  with  $|V(R')| = r'$  such that  $2r_0 + 1 \leq r'_0 \leq r' \leq r_1$  and with minimum degree at least  $(\frac{1}{2} + \gamma - d - \varepsilon')r' \geq (\frac{1}{2} + \frac{\gamma}{2})r'$  by (23). If  $r'$  is odd, then set  $V_0 := V'_0 \dot{\cup} V'_{r'}$  and  $r := (r' - 1)/2$ , otherwise set  $V_0 := V'_0$  and  $r := r'/2$ . Clearly  $r_0 \leq r \leq r_1$ , the graph  $R := R'[2r]$  still has minimum degree at least  $(\frac{1}{2} + \frac{\gamma}{3})2r$  and  $|V_0| \leq \varepsilon'n + (n/r'_0) \leq (\eta/5)n$  by the choice of  $r'_0$  and  $\varepsilon'$ . It follows from Theorem 2 applied with  $\Delta = 3$  and  $\gamma/2$  that  $R$  contains a copy of the ladder  $R_r^*$  on  $2r$  vertices ( $R_r^*$  has bandwidth  $2 \leq \beta \cdot 2r$  by the choice of  $r'_0$  in (22)). Hence we can rename the vertices of the graph  $R = R'[2r]$  with  $u_1, v_1, \dots, u_r, v_r$  according to the spanning copy of  $R_r^*$ . This naturally defines an equipartite mapping  $f$  from  $V \setminus V_0$  to the vertices of the ladder  $R_r^*$ , where  $f$  maps all vertices in some cluster  $V_i$  with  $i \in [2r]$  to a vertex  $u_{i'}$  or  $v_{i'}$  of  $R_r^*$  for some index  $i' \in [r]$ . We will show that subdividing the clusters  $f^{-1}(x)$  for all  $x \in V(R_r^*)$  will give the desired mapping  $g$ .




FIGURE 3. Cutting off a set of balancing clusters from  $f^{-1}(u_i)$  and  $f^{-1}(w_i)$ . These clusters build  $p$ -dense pairs (thanks to the triangle  $u_i v_i w_i$  in  $R$ ) in the form of a  $C_5$ .

We will now construct the balancing clusters  $B_{i,j}$  and  $B'_{i,j}$  with  $i \in [r], j \in [2t]$  and afterwards turn to the connecting clusters  $C_{i,j}$  and  $C'_{i,j}$  and big clusters  $U_i$  and  $V_i$  with  $i \in [r], j \in [2t]$ .

Notice that  $\delta(R) \geq (\frac{1}{2} + \frac{\gamma}{3})2r$  implies that every edge  $u_i v_i$  of  $R_r^* \subseteq R$  is contained in more than  $\gamma r$  triangles in  $R$ . Therefore, we can choose vertices  $w_i$  of  $R$  for all  $i \in [r]$  such that  $u_i v_i w_i$  forms a triangle in  $R$  and no vertex of  $R$  serves as  $w_i$  more than  $2/\gamma$  times. We continue by choosing in cluster  $f^{-1}(u_i)$  arbitrary disjoint vertex sets  $B_{i,1}, \dots, B_{i,t}, B'_{i,t+1}, \dots, B'_{i,2t}$ , of size  $\eta'n/(2r)$  each, for all  $i \in [r]$ . We will show below that  $f^{-1}(u_i)$  is large enough so that these sets can be chosen. We then remove all vertices in these sets from  $f^{-1}(u_i)$ . Similarly, we choose in cluster  $f^{-1}(w_i)$  arbitrary disjoint vertex sets  $B_{i,t+1}, \dots, B_{i,2t}, B'_{i,1}, \dots, B'_{i,t}$ , of size  $\eta'n/(2r)$  each, for all  $i \in [r]$ . We also remove these sets from  $f^{-1}(w_i)$ . Observe that this construction asserts the following property. For all  $i \in [r]$  and  $j, j', j'', j''' \in [t]$  each of the pairs  $(f^{-1}(v_i), B_{i,j}), (B_{i,j}, B'_{i,j'}), (B'_{i,j'}, B'_{i,t+j''}), (B'_{i,t+j''}, B_{i,t+j'''}), (B_{i,t+j'''}, f^{-1}(v_i))$  is a sub-pair of a  $p$ -dense

pair corresponding to an edge of  $R[\{u_i, v_i, w_i\}]$  (see Figure 3). Accordingly this is a sequence of  $p$ -dense pairs in the form of a  $C_5$ , as needed for the balancing clusters in view of condition (G3) (see also Figure 1). Hence we call the sets  $B_{i,j}$  and  $B'_{i,j}$  with  $i \in [r]$ ,  $j \in [2t]$  balancing clusters from now on and claim that they have the required properties. This claim will be verified below.

We now turn to the construction of the connecting clusters and big clusters. Recall that we already removed balancing clusters from all clusters  $f^{-1}(u_i)$  and possibly from some clusters  $f^{-1}(v_i)$  (because  $v_i$  might have served as  $w_{i'}$ ) with  $i \in [r]$ . For each  $i \in [r]$  we arbitrarily partition the remaining vertices of cluster  $f^{-1}(u_i)$  into sets  $C_{i,1} \dot{\cup} \dots \dot{\cup} C_{i,2t} \dot{\cup} U_i$  and the remaining vertices of cluster  $f^{-1}(v_i)$  into sets  $C'_{i,1} \dot{\cup} \dots \dot{\cup} C'_{i,2t} \dot{\cup} V_i$  such that  $|C_{i,j}|, |C'_{i,j}| = \eta'n/(2r)$  for all  $i \in [r]$ ,  $j \in [2t]$ . This gives us the connecting and the big clusters and we claim that also these clusters have the required properties. Observe, again, that for all  $i \in [r]$ ,  $i' \in \{i-1, i, i+1\} \setminus \{0\}$ ,  $j, j' \in [2t]$  each of the pairs  $(U_i, V_i)$ ,  $(C_{i',j}, V_i)$ , and  $(C_{i,j}, C'_{i',j'})$  is a sub-pair of a  $p$ -dense pair corresponding to an edge of  $R_r^*$  (see Figure 4).




FIGURE 4. Partitioning the remaining vertices of cluster  $f^{-1}(u_i)$  and  $f^{-1}(v_i)$  into sets  $C_{i,1} \dot{\cup} C_{i,2} \dot{\cup} U_i$  and  $C'_{i,1} \dot{\cup} C'_{i,2} \dot{\cup} V_i$  (for the special case  $t = 1$ ). These clusters form  $p$ -dense pairs (thanks to the ladder  $R_r^*$  in  $R$ ) as indicated by the edges.

We will now show that the balancing clusters, connecting clusters and big clusters satisfy conditions (G1)–(G3). Note that condition (G2) concerning the sizes of the connecting and balancing clusters is satisfied by construction. To determine the sizes of the big clusters observe that from each cluster  $V'_j$  with  $j \in [2r]$  vertices for at most  $2t \cdot 2/\gamma$  balancing clusters were removed. In addition, at most  $2t$  connecting clusters were split off from  $V'_j$ . Since  $|V \setminus V_0| \geq (1 - \eta/5)n$  we get


$$|V_i|, |U_i| \geq \left(1 - \frac{\eta}{5}\right) \frac{n}{2r} - \left(\frac{4}{\gamma} + 2\right) t \cdot \eta' \frac{n}{2r} \geq (1 - \eta) \frac{n}{2r}$$

by (21). This is condition (G1). It remains to verify condition (G3). It can easily be checked that for all  $xy \in E(R_{r,t})$  the corresponding pair  $(g^{-1}(x), g^{-1}(y))$  is a sub-pair of some cluster pair  $(f^{-1}(x'), f^{-1}(y'))$  with  $x'y' \in E(R)$  by construction. In addition, all big, connecting, and balancing clusters are of size at least  $\eta'n/(2r)$ . Hence we have  $|g^{-1}(x)| \geq \eta'|f^{-1}(x')|$  and  $|g^{-1}(y)| \geq \eta'|f^{-1}(y')|$ . We conclude from Proposition 6 that  $(g^{-1}(x), g^{-1}(y))$  is  $(\varepsilon, d, p)$ -dense since  $\varepsilon'/\eta' \leq \varepsilon$  by (23). This finishes the verification of (G3).  $\square$

## 10. A PARTITION OF $H$

Hajnal and Szemerédi determined the minimum degree that forces a certain number of vertex disjoint  $K_r$  copies in  $G$ . In addition their result guarantees that the remaining vertices can be covered by copies of  $K_{r-1}$ . Another way to express this, which actually resembles the original formulation, is obtained by considering the complement  $\bar{G}$  of  $G$  and its maximum degree. Then, so the theorem asserts, the graph  $\bar{G}$  contains a certain number of vertex disjoint independent sets of almost equal sizes. In other words,  $\bar{G}$  admits a vertex colouring such that the sizes of the colour classes differ by at most 1. Such a colouring is also called *equitable colouring*.

**Theorem 22** (Hajnal & Szemerédi [18]). *Let  $\bar{G}$  be a graph on  $n$  vertices with maximum degree  $\Delta(\bar{G}) \leq \Delta$ . Then there is an equitable vertex colouring of  $G$  with  $\Delta + 1$  colours.*  $\square$

FIGURE 5. The graph  $\bar{R}$  in Proposition 23.

In the proof of Lemma 11 that we shall present in this section we will use this theorem in order to guarantee property (H3). This will be the very last step in the proof, however. First, we need to take care of the remaining properties.

Before we start, let us agree on some terminology that will turn out to be useful in the proof of Lemma 11. When defining a homomorphism  $h$  from a graph  $H$  to a graph  $R$ , we write  $h(S) := z$  for a set  $S$  of vertices in  $H$  and a vertex  $z$  in  $R$  to say that all vertices from  $S$  are mapped to  $z$ . Recall that we have a bandwidth hypothesis on  $H$ . Consider an ordering of the vertices of  $H$  achieving its bandwidth. Then we can deal with the vertices of  $H$  in this order. In particular, we can refer to vertices as the *first* or *last* vertices in some set, meaning that they are the vertices with the smallest or largest label from this set.

We start with the following proposition.

**Proposition 23.** *Let  $\bar{R}$  be the following graph with six vertices and six edges:*

$$\bar{R} := (\{z^0, z^1, \dots, z^5\}, \{z^0 z^1, z^1 z^2, z^2 z^3, z^3 z^4, z^4 z^5, z^5 z^1\}),$$

*see Figure 5 for a picture of  $\bar{R}$ . For every real  $\bar{\eta} > 0$  there exists a real  $\bar{\beta} > 0$  such that the following holds: Consider an arbitrary bipartite graph  $\bar{H}$  with  $\bar{m}$  vertices, colour classes  $Z^0$  and  $Z^1$ , and  $\text{bw}(\bar{H}) \leq \bar{\beta}\bar{m}$  and denote by  $T$  the union of the first  $\bar{\beta}\bar{m}$  vertices and the last  $\bar{\beta}\bar{m}$  vertices of  $\bar{H}$ . Then there exists a homomorphism  $\bar{h}: V(\bar{H}) \rightarrow V(\bar{R})$  from  $\bar{H}$  to  $\bar{R}$  such that for all  $j \in \{0, 1\}$  and all  $k \in [2, 5]$*

$$\frac{\bar{m}}{2} - 5\bar{\eta}\bar{m} \leq |\bar{h}^{-1}(z^j)| \leq \frac{\bar{m}}{2} + \bar{\eta}\bar{m}, \quad (24)$$

$$|\bar{h}^{-1}(z^k)| \leq \bar{\eta}\bar{m}, \quad (25)$$

$$h(T \cap Z^j) = z^j. \quad (26)$$

Roughly speaking, Proposition 23 shows that we can find a homomorphism from a bipartite graph  $\bar{H}$  to a graph  $\bar{R}$  which consists of an edge  $z^0 z^1$  which has an attached 5-cycle in such a way that most of the vertices of  $\bar{H}$  are mapped about evenly to the vertices  $z^0$  and  $z^1$ . If we knew that the colour classes of  $\bar{H}$  were of almost equal size, then this would be a trivial task, but since this is not guaranteed, we will have to make use of the additional vertices  $z^2, \dots, z^5$ .

*Proof of Proposition 23.* Given  $\bar{\eta}$ , choose an integer  $\ell \geq 6$  and a real  $\bar{\beta} > 0$  such that

$$\frac{5}{\ell} < \bar{\eta} \quad \text{and} \quad \bar{\beta} := \frac{1}{\ell^2}. \quad (27)$$

For the sake of a simpler exposition we assume that  $\bar{m}/\ell$  and  $\bar{\beta}\bar{m}$  are integers. Now consider a graph  $\bar{H}$  as given in the statement of the proposition. Partition  $V(\bar{H})$  along the ordering induced by the bandwidth labelling into sets  $\bar{W}_1, \dots, \bar{W}_\ell$  of sizes  $|\bar{W}_i| = \bar{m}/\ell$  for  $i \in [\ell]$ . For each  $\bar{W}_i$ , consider its last  $5\bar{\beta}\bar{m}$  vertices and partition them into sets  $X_{i,1}, \dots, X_{i,5}$  of size  $|X_{i,k}| = \bar{\beta}\bar{m}$ . For  $i \in [\ell]$  let

$$W_i := \bar{W}_i \setminus (X_{i,1} \cup \dots \cup X_{i,5}), \quad W := \bigcup_{i=1}^{\ell} W_i,$$



and note that

$$L := |W_i| = \frac{\bar{m}}{\ell} - 5\bar{\beta}\bar{m} \stackrel{(27)}{=} \left(\frac{1}{\ell} - \frac{5}{\ell^2}\right) \bar{m} \geq \frac{1}{\ell^2} \bar{m} \stackrel{(27)}{=} \bar{\beta}\bar{m}.$$

For  $i \in [\ell]$ ,  $j \in \{0, 1\}$ , and  $1 \leq k \leq 5$  let

$$W_i^j := W_i \cap Z^j, \quad X_{i,k}^j := X_{i,k} \cap Z^j.$$

Thanks to the fact that  $\text{bw}(\bar{H}) \leq \bar{\beta}\bar{m}$ , we know that there are no edges between  $W_i$  and  $W_{i'}$  for  $i \neq i' \in [\ell]$ . In a first round, for each  $i \in [\ell]$  we will either map all vertices from  $W_i^j$  to  $z^j$  for both  $j \in \{0, 1\}$  (call such a mapping a *normal* embedding of  $W_i$ ) or we map all vertices from  $W_i^j$  to  $z^{1-j}$  for both  $j \in \{0, 1\}$  (call this an *inverted* embedding). We will do this in such a way that the difference between the number of vertices that get sent to  $z^0$  and the number of those that get sent to  $z^1$  is as small as possible. Since  $|W_i| \leq L$  the difference is therefore at most  $L$ . If, in addition, we guarantee that both  $W_1$  and  $W_\ell$  receive a normal embedding, it is at most  $2L$ . So, to summarize and to describe the mapping more precisely: there exist integers  $\varphi_i \in \{0, 1\}$  for all  $i \in [\ell]$  such that  $\varphi_1 = 0 = \varphi_\ell$  and the function  $h : W \rightarrow \{z^0, z^1\}$  defined by

$$h(W_i^j) := \begin{cases} z^j & \text{if } \varphi_i = 0, \\ z^{1-j} & \text{if } \varphi_i = 1, \end{cases}$$

is a homomorphism from  $\bar{H}[W]$  to  $\bar{R}[\{z^0, z^1\}]$ , satisfying that for both  $j \in \{0, 1\}$

$$\begin{aligned} |h^{-1}(z^j)| &\leq \frac{\ell L}{2} + 2L = \left(\frac{\ell}{2} + 2\right) \frac{\bar{m}}{\ell} - \left(\frac{\ell}{2} + 2\right) 5\bar{\beta}\bar{m} \\ &\stackrel{(27)}{=} \frac{\bar{m}}{2} + \bar{m} \left(\frac{2}{\ell} - \frac{5}{2\ell} - \frac{10}{\ell^2}\right) \leq \frac{\bar{m}}{2}. \end{aligned} \tag{28}$$

In the second round we extend this homomorphism to the vertices in the classes  $X_{i,k}$ . Recall that these vertices are by definition situated after those in  $W_i$  and before those in  $W_{i+1}$ . The idea for the extension is simple. If  $W_i$  and  $W_{i+1}$  have been embedded in the same way by  $h$  (either both normal or both inverted), then we map all the vertices from all  $X_{i,k}$  to  $z^0$  and  $z^1$  accordingly. If they have been embedded in different ways (one normal and one inverted), then we walk around the 5-cycle  $z^1, \dots, z^5, z^1$  to switch colour classes.

Here is the precise definition. Consider an arbitrary  $i \in [\ell]$ . Since  $h(W_i^0)$  and  $h(W_i^1)$  are already defined, choose (and fix)  $j \in \{0, 1\}$  in such a way that  $h(W_i^j) = z^1$ . Note that this implies that  $h(W_i^{1-j}) = z^0$ . Now define  $h_i : \bigcup_{k=0}^5 X_{i,k} \rightarrow \bigcup_{k=1}^5 \{z^k\}$  as follows:

Suppose first that  $\varphi_i = \varphi_{i+1}$ . Observe that in this case we must also have  $h(W_{i+1}^j) = z^1$  and  $h(W_{i+1}^{1-j}) = z^0$ . So we can happily define for all  $k \in [5]$

$$h_i(X_{i,k}^j) = z^1 \quad \text{and} \quad h_i(X_{i,k}^{1-j}) = z^0.$$

Now suppose that  $\varphi_i \neq \varphi_{i+1}$ . Since we are still assuming that  $j$  is such that  $h(W_i^j) = z^1$  and thus  $h(W_i^{1-j}) = z^0$ , the fact that  $\varphi_i \neq \varphi_{i+1}$  implies that  $h(W_{i+1}^j) = z^0$  and  $h(W_{i+1}^{1-j}) = z^1$ . In this case we define  $h_i$  as follows:

|                           |                                  |                                  |                                  |                                  |                                  |                               |
|---------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|-------------------------------|
| $h(W_i^{1-j})$<br>$= z^0$ | $h_i(X_{i,1}^{1-j})$<br>$:= z^2$ | $h_i(X_{i,2}^{1-j})$<br>$:= z^2$ | $h_i(X_{i,3}^{1-j})$<br>$:= z^4$ | $h_i(X_{i,4}^{1-j})$<br>$:= z^4$ | $h_i(X_{i,5}^{1-j})$<br>$:= z^1$ | $h(W_{i+1}^{1-j})$<br>$= z^1$ |
| $h(W_i^j)$<br>$= z^1$     | $h_i(X_{i,1}^j)$<br>$:= z^1$     | $h_i(X_{i,2}^j)$<br>$:= z^3$     | $h_i(X_{i,3}^j)$<br>$:= z^3$     | $h_i(X_{i,4}^j)$<br>$:= z^5$     | $h_i(X_{i,5}^j)$<br>$:= z^5$     | $h(W_{i+1}^j)$<br>$= z^0$     |

Finally, we set  $\bar{h} : V(\bar{H}) \rightarrow V(\bar{R})$  by letting  $\bar{h}(x) := h(x)$  if  $x \in W_i$  for some  $i \in [\ell]$  and  $\bar{h}(x) := h_i(x)$  if  $x \in X_{i,k}$  for some  $i \in [\ell]$  and  $k \in [5]$ .

In order to verify that this is a homomorphism from  $\bar{H}$  to the sets  $\bar{R}$ , we first let

$$X_{i,0}^0 := W_i^0, X_{i,0}^1 := W_i^1, X_{i,6}^0 := W_{i+1}^0, X_{i,6}^1 := W_{i+1}^1.$$

Using this notation, it is clear that any edge  $xx'$  in  $\bar{H}[W_i \cup \bigcup_{k=1}^5 X_{i,k} \cup W_{i+1}]$  with  $x \in Z^j$  and  $x' \in Z^{1-j}$  is of the form

$$xx' \in (X_{i,k}^j \times X_{i,k}^{1-j}) \cup (X_{i,k}^j \times X_{i,k+1}^{1-j}) \cup (X_{i,k+1}^j \times X_{i,k}^{1-j})$$

for some  $k \in [0, 6]$ . It is therefore easy to check in the above table that  $\bar{h}$  maps  $xx'$  to an edge of  $R$ .

We conclude the proof by showing that the cardinalities of the preimages of the vertices in  $R$  match the required sizes. In the second round we mapped a total of

$$\ell \cdot 5\bar{\beta}\bar{m} \stackrel{(27)}{=} \frac{5}{\ell}\bar{m} \stackrel{(27)}{\leq} \bar{\eta}\bar{m}$$

additional vertices from  $\bar{H}$  to the vertices of  $\bar{R}$ , which guarantees that

$$|\bar{h}^{-1}(z^j)| \stackrel{(28)}{\leq} \frac{\bar{m}}{2} + \bar{\eta}\bar{m} \text{ for all } j \in \{0, 1\}, \quad |\bar{h}^{-1}(z^k)| \leq \bar{\eta}\bar{m} \text{ for all } k \in [2, 5].$$

Finally, the lower bound in (24) immediately follows from the upper bounds:

$$|\bar{h}^{-1}(z^j)| \geq \bar{m} - |\bar{h}^{-1}(z^{1-j})| - \sum_{k=2}^5 |\bar{h}^{-1}(z^k)| \geq \frac{\bar{m}}{2} - 5\bar{\eta}\bar{m}.$$

□

We remark that Proposition 23 (and thus Lemma 11) would remain true if we replaced the 5-cycle in  $\bar{R}$  by a 3-cycle. However, we need the properties of the 5-cycle in the proof of the main theorem. Now we will prove Lemma 11.

*Proof of Lemma 11.* Given the integer  $\Delta$ , set  $t := (\Delta + 1)^3(\Delta^3 + 1)$ . Given a real  $0 < \eta < 1$  and integers  $m$  and  $r$ , set  $\bar{\eta} := \eta/20 < 1/20$  and apply Proposition 23 to obtain a real  $\bar{\beta} > 0$ . Choose  $\beta > 0$  sufficiently small so that all the inequalities

$$\frac{1}{r} - 4\beta \geq \beta/\bar{\beta}, \quad 4\beta r \leq \frac{\eta}{20r}, \quad 16\Delta\beta r \leq \eta \left( \frac{1}{r} - 4\beta \right) \left( \frac{1}{2} - 5\bar{\eta} \right) \quad (29)$$

hold. Again, we assume that  $m/r$  and  $\beta m$  are integers.

Next we consider the spin graph  $R_{r,t}$  with  $t = 1$ , i.e., let  $R := R_{r,1}$ . For the sake of simpler reference, we will change the names of its vertices as follows: For all  $i \in [r]$  we set (see Figure 6)

$$\begin{aligned} z_i^0 &:= u_i, z_i^1 := v_i, z_i^2 := b_{i,1}, z_i^3 := b'_{i,1}, z_i^4 := b'_{i,2}, z_i^5 := b_{i,2}, \\ q_i^2 &:= c_{i,1}, q_i^3 := c'_{i,1}, q_i^4 := c_{i,2}, q_i^5 := c'_{i,2}. \end{aligned}$$

Note that for every  $i \in [r]$  the graph  $R[\{z_i^0, \dots, z_i^5\}]$  is isomorphic to the graph  $\bar{R}$  defined in Proposition 23.

Partition  $V(H)$  along the ordering (induced by the bandwidth labelling) into sets  $\bar{S}_1, \dots, \bar{S}_r$  of sizes  $|\bar{S}_i| = m/r$  for  $i \in [r]$ .

Define sets  $T_{i,k}$  for  $i \in [r]$  and  $k \in [0, 5]$  with  $|T_{i,k}| = \beta m$  such that  $T_{i,0} \cup \dots \cup T_{i,4}$  contain the last  $5\beta m$  vertices of  $\bar{S}_i$  and  $T_{i,5}$  the first  $\beta m$  vertices of  $\bar{S}_{i+1}$  (according to the ordering). Set  $S_i := \bar{S}_i \setminus (T_{i,1} \cup \dots \cup T_{i,4})$  and observe that this implies that  $T_{i,0}$  is the set of the last  $\beta m$  vertices of  $S_i$  and  $T_{i,5}$  is the set of the first  $\beta m$  vertices in  $\bar{S}_{i+1}$ . Set

$$\bar{m} := |S_i| = (m/r) - 4\beta m = \left( \frac{1}{r} - 4\beta \right) m \stackrel{(29)}{\geq} \beta m / \bar{\beta}, \quad \text{thus} \quad \bar{\beta}\bar{m} \geq \beta m. \quad (30)$$

Denote by  $Z^0$  and  $Z^1$  the two colour classes of the bipartite graph  $H$ . For  $i \in [\ell]$ ,  $k \in [0, 5]$  and  $j \in [0, 1]$  let

$$S_i^j := S_i \cap Z^j, \quad T_{i,k}^j := T_{i,k} \cap Z^j.$$

Now for each  $i \in [r]$  apply Proposition 23 to  $\bar{H}_i := H[S_i]$  and  $\bar{R}_i := R[\{z_i^0, \dots, z_i^5\}]$ . Observe that

$$\text{bw}(\bar{H}_i) \leq \text{bw}(H) \leq \beta m \stackrel{(30)}{\leq} \bar{\beta}\bar{m},$$




FIGURE 6. The subgraph  $R[\{z_i^0, \dots, z_i^5, q_i^4, q_i^5, q_{i+1}^2, q_{i+1}^3, z_{i+1}^0, \dots, z_{i+1}^5\}]$  of  $R_{r,1}$  in the proof of Lemma 11.

so we obtain a homomorphism  $\bar{h}_i : S_i \rightarrow \{z_i^0, \dots, z_i^5\}$  of  $\bar{H}_i$  to  $\bar{R}_i$ . Combining these yields a homomorphism

$$\begin{aligned} \bar{h} : \bigcup_{i=1}^r S_i &\rightarrow \bigcup_{i=1}^r \{z_i^0, \dots, z_i^5\}, \\ \text{from } H[\bigcup_{i=1}^r S_i] &\text{ to } R[\bigcup_{i=1}^r \{z_i^0, \dots, z_i^5\}] \end{aligned}$$

with the property that for every  $i \in [r]$ ,  $j \in [0, 1]$  and  $k \in [2, 5]$

$$\begin{aligned} \frac{\bar{m}}{2} - 5\bar{\eta}\bar{m} &\stackrel{(24)}{\leq} |\bar{h}^{-1}(z_i^j)| \stackrel{(24)}{\leq} \frac{\bar{m}}{2} + \bar{\eta}\bar{m} \leq \left(1 + \frac{\eta}{10}\right) \frac{m}{2r} \quad \text{and} \\ |\bar{h}^{-1}(z_i^k)| &\stackrel{(25)}{\leq} \bar{\eta}\bar{m} \leq \frac{\eta}{10} \frac{m}{2r}. \end{aligned}$$

Thanks to (30), we know that  $\bar{\beta}\bar{m} \geq \beta m$ , and therefore applying the information from (26) in Proposition 23 yields that for all  $i \in [r]$  and  $j \in [0, 1]$

$$\bar{h}(T_{i,0}^j) = z_i^j \quad \text{and} \quad \bar{h}(T_{i,5}^j) = z_{i+1}^j.$$

In the second round, our task is to extend this homomorphism to the vertices in  $\bar{S}_i \setminus S_i$  by defining a function

$$h_i : T_{i,1} \cup \dots \cup T_{i,4} \rightarrow \{z_i^1, q_i^4, q_i^5, q_{i+1}^2, q_{i+1}^3, z_{i+1}^1\}$$

for each  $i \in [r]$  as follows:

|                              |                           |                           |                               |                               |                                  |
|------------------------------|---------------------------|---------------------------|-------------------------------|-------------------------------|----------------------------------|
| $\bar{h}(T_{i,0}^0) = z_i^0$ | $h_i(T_{i,1}^0) := q_i^4$ | $h_i(T_{i,2}^0) := q_i^4$ | $h_i(T_{i,3}^0) := q_{i+1}^2$ | $h_i(T_{i,4}^0) := q_{i+1}^3$ | $\bar{h}(T_{i,5}^0) = z_{i+1}^0$ |
| $\bar{h}(T_{i,0}^1) = z_i^1$ | $h_i(T_{i,1}^1) := z_i^1$ | $h_i(T_{i,2}^1) := q_i^5$ | $h_i(T_{i,3}^1) := q_{i+1}^3$ | $h_i(T_{i,4}^1) := z_{i+1}^1$ | $\bar{h}(T_{i,5}^1) = z_{i+1}^1$ |

Now set  $h(x) := \bar{h}(x)$  if  $x \in S_i$  for some  $i \in [r]$  and  $h(x) := h_i(x)$  if  $x \in T_{i,k}$  for some  $i \in [r]$  and  $k \in [4]$ .

Let us verify that  $h$  is a homomorphism from  $H$  to  $R$ . For edges  $xx'$  with both endpoints inside a set  $S_i$  we do not need to check anything because here  $h(x) = \bar{h}(x)$  and  $h(x') = \bar{h}(x')$  and we know from Proposition 23 that  $\bar{h}$  is a homomorphism. Due to the bandwidth condition  $\text{bw}(H) \leq \beta m$ , any other edge  $xx'$  with  $x \in Z^0$  and  $x' \in Z^1$  is of the form

$$xx' \in (T_{i,k}^0 \times T_{i,k}^1) \cup (T_{i,k}^0 \times T_{i,k+1}^1) \cup (T_{i,k+1}^0 \times T_{i,k}^1)$$

for some  $i \in [r]$  and  $0 \leq k, k+1 \leq 5$ . It is therefore easy to check in the above table that  $h$  maps  $xx'$  to an edge of  $R$ .

What can we say about the cardinalities of the preimages? In the second round we have mapped  $4\beta mr$  additional vertices from  $H$  to vertices in  $R$ , hence for any vertex  $z$  in  $R$  with  $z \notin \{z_i^0, z_i^1\}$ ,  $i \in [\ell]$  we have

$$|h_i^{-1}(z)| \leq 4\beta mr \stackrel{(29)}{\leq} \frac{\eta}{10} \frac{m}{2r}, \quad (31)$$

and therefore the required upper bounds immediately follow from (10).

At this point we have found a homomorphism  $h$  from  $H$  to  $R = R_{r,1}$  of which we know that it satisfies properties (H1) and (H2).

So far we have been working with the graph  $R = R_{r,1}$ , and therefore we know which vertices have been mapped to  $u_i = z_i^0$  and  $v_i = z_i^1$ :

$$\widetilde{U}_i := h^{-1}(u_i) = h^{-1}(z_i^0) \quad \text{and} \quad \widetilde{V}_i := h^{-1}(v_i) = h^{-1}(z_i^1).$$

Moreover for  $i \in [r]$  and  $k \in [2, 5]$  set

$$Z_i^k := h^{-1}(z_i^k) \quad \text{and} \quad Q_i^k := h^{-1}(q_i^k)$$

Let us deal with property (H5) next. By definition, a vertex in  $\widetilde{X}_i \subseteq \widetilde{V}_i$  must have at least one neighbour in  $Q_i^2$  or  $Q_i^4$  or  $Z_i^2$  or  $Z_i^5$ . We know from (31) that the two latter sets contain at most  $4\beta mr$  vertices each, and each of their vertices has at most  $\Delta$  neighbours. Thus

$$\begin{aligned} |\widetilde{X}_i| &\leq \Delta \cdot 16\beta mr \stackrel{(29)}{\leq} \eta \left( \frac{1}{r} - 4\beta \right) \left( \frac{1}{2} - 5\bar{\eta} \right) m \stackrel{(30)}{=} \eta \bar{m} \left( \frac{1}{2} - 5\bar{\eta} \right) \\ &\stackrel{(10)}{\leq} \eta |\bar{h}^{-1}(z_i^1)| \leq \eta |h^{-1}(z_i^1)| = \eta |\widetilde{V}_i|, \end{aligned}$$

which shows that (H5) is also satisfied.

Next we would like to split up the sets  $Z_i^k$  and  $Q_i^k$  for  $i \in [r]$  and  $k \in [2, 5]$  into smaller sets in order to meet the additional requirements (H3) and (H4). This means that we need to partition them further into sets of vertices which have no path of length 1, 2, or 3 between them and which have the same degree into certain sets.

To achieve this, first denote by  $H^3$  the 3rd power of  $H$ . Then an upper bound on the maximum degree of  $H^3$  is obviously given by

$$\Delta + \Delta(\Delta - 1) + \Delta(\Delta - 1)(\Delta - 1) \leq \Delta^3.$$

Hence  $H^3$  has a vertex colouring  $c : V(H) \rightarrow \mathbb{N}$  with at most  $\Delta^3 + 1$  colours. Notice that a set of vertices that receives the same colour by  $c$  forms a 3-independent set in  $H$ . To formalize this argument, we define a ‘fingerprint’ function

$$f : \bigcup_{i=1}^r \bigcup_{k=2}^5 (Z_i^k \cup Q_i^k) \rightarrow [0, \Delta] \times [0, \Delta] \times [0, \Delta] \times [\Delta^3 + 1]$$

as follows:

$$f(y) := \begin{cases} \left( \deg_{\widetilde{V}_i}(y), \deg_{Q_i^2 \cup Q_i^4}(y), \deg_{Z_i^2}(y), c(y) \right) & \text{if } y \in \left( \bigcup_{k=2}^5 (Q_i^k \cup Z_i^k) \right) \setminus Z_i^4, \\ \left( \deg_{\widetilde{V}_i}(y), \deg_{Q_i^2 \cup Q_i^4}(y), \deg_{Z_i^3 \cup Z_i^5}(y), c(y) \right) & \text{if } y \in Z_i^4, \end{cases}$$

for some  $i \in [r]$ .

Recall that we defined  $t := (\Delta + 1)^3(\Delta^3 + 1)$ , so let us identify the codomain of  $f$  with the set  $[t]$ . Now for  $i \in [r]$  and  $j \in [t]$  we set

$$\begin{aligned} \widetilde{B}_{i,j} &:= Z_i^2 \cap f^{-1}(j), & \widetilde{B}_{i,t+j} &:= Z_i^5 \cap f^{-1}(j) \\ \widetilde{B}'_{i,j} &:= Z_i^3 \cap f^{-1}(j), & \widetilde{B}'_{i,t+j} &:= Z_i^4 \cap f^{-1}(j) \\ \widetilde{C}_{i,j} &:= Q_i^2 \cap f^{-1}(j), & \widetilde{C}_{i,t+j} &:= Q_i^4 \cap f^{-1}(j) \\ \widetilde{C}'_{i,j} &:= Q_i^3 \cap f^{-1}(j), & \widetilde{C}'_{i,t+j} &:= Q_i^5 \cap f^{-1}(j). \end{aligned}$$

Observe, for example, that for  $y \in \tilde{B}_{i,j}$  the third component of  $f(y)$  is exactly equal to  $\deg_{L(i,j)}(y)$ . Now, for any

$$yy' \in \binom{\tilde{C}_{i,j}}{2} \cup \binom{\tilde{B}_{i,j}}{2} \cup \binom{\tilde{C}'_{i,j}}{2} \cup \binom{\tilde{B}'_{i,j}}{2},$$

we have  $f(y) = j = f(y')$  and hence any of the parameters required in (H3) and (H4) have the same value for  $y$  and  $y'$ .

The only thing missing before the proof of Lemma 11 is complete is that we need to guarantee that every  $y \in Z_i^2 \cup Z_i^5 \cup Q_i^2 \cup Q_i^4$  has at most  $\Delta - 1$  neighbours in  $\tilde{V}_i$ , as required in the first line of (H4). If a vertex  $y$  does not satisfy this, it must have *all* its  $\Delta$  neighbours in  $\tilde{V}_i$ . Since by definition of  $\tilde{V}_i$  these neighbours have been mapped to  $z_i^1$ , we can map  $y$  to  $z_i^0$  (instead of mapping it to  $z_i^2, z_i^5, q_i^2$  or  $q_i^4$ ).

Even if, in this way, all of the vertices in  $Z_i^2 \cup Z_i^5 \cup Q_i^2 \cup Q_i^4$  would have to be mapped to  $z_i^0$ , (31) assures us that these are at most  $4\frac{\eta}{10}\frac{m}{2r}$  vertices. Since by (10) at most  $(1 + \frac{\eta}{10})\frac{m}{2r}$  have already been mapped to  $z_i^0$  in the first round and by (31) at most  $\frac{\eta}{10}\frac{m}{2r}$  in the second round, this does not violate the upper bound in (H1).  $\square$

## 11. THE CONSTRAINED BLOW-UP LEMMA

As explained earlier, the proof of the constrained blow-up lemma uses techniques developed in [4, 25] adapted to our setting. In fact, the proof we present here follows the embedding strategy used in the proof of [4, Theorem 1.5]. This strategy is roughly as follows. Assume we want to embed the bipartite graph  $H$  on vertex set  $\tilde{U} \dot{\cup} \tilde{V}$  into the host graph  $G$  on vertex set  $U \dot{\cup} V$ . Then we consider injective mappings  $f: \tilde{V} \rightarrow V$ , and try to find one that can be extended to  $\tilde{U}$  such that the resulting mapping is an embedding of  $H$  into  $G$ . For determining whether a particular mapping  $f$  can be extended in this way we shall construct an auxiliary bipartite graph  $B_f$ , the so-called candidate graph (see Definition 24), which contains a matching covering one of its partition classes if and only if  $f$  can be extended. Accordingly, our goal will be to check whether  $B_f$  contains such a matching  $M$  which we will do by appealing to Hall's condition. On page 27 we will explain the details of this part of the proof, determine necessary conditions for the application of Hall's theorem, and collect them in form of a matching lemma (Lemma 31). It will then remain to show that there is a mapping  $f$  such that  $B_f$  satisfies the conditions of this matching lemma. This will require most of the work. The idea here is as follows.

We will show that mappings  $f$  usually have the necessary properties as long as they do not map neighbourhoods  $N_H(\tilde{u}) \subseteq \tilde{V}$  of vertices in  $\tilde{u} \in \tilde{U}$  to certain “bad” spots in  $V$ . The existence of (many) mappings that avoid these “bad” spots is verified with the help of a hypergraph packing lemma (Lemma 29). This lemma states that half of all possible mappings  $f$  avoid *almost all* “bad” spots and can easily be turned into mappings  $f'$  avoiding *all* “bad” spots with the help of so-called switchings.

**11.1. Candidate graphs.** If we have injective mappings  $f: \tilde{V} \rightarrow V$  as described in the previous paragraph we would like to decide whether  $f$  can be extended to an embedding of  $H$  into  $G$ . Observe that in such an embedding each vertex  $\tilde{u} \in \tilde{U}$  has to be embedded to a vertex  $u \in U$  such that the following holds. The neighbourhood  $N_H(\tilde{u})$  has its image  $f(N_H(\tilde{u}))$  in the set  $N_G(u)$ . Determining which vertices  $u$  are “candidates” for the embedding of  $\tilde{u}$  in this sense gives rise to the following bipartite graph.

**Definition 24** (candidate graph). *Let  $H$  and  $G$  be bipartite graphs on vertex sets  $\tilde{U} \dot{\cup} \tilde{V}$  and  $U \dot{\cup} V$ , respectively. For an injective function  $f: \tilde{V} \rightarrow V$  we say that a vertex  $u \in U$  is an  $f$ -candidate for  $\tilde{u} \in \tilde{U}$  if and only if  $f(N_H(\tilde{u})) \subseteq N_G(u)$ .*

*The candidate graph  $B_f(H, G) := (\tilde{U} \dot{\cup} U, E_f)$  for  $f$  is the bipartite graph with edge set*

$$E_f := \left\{ \tilde{u}u \in \tilde{U} \times U : u \text{ is an } f\text{-candidate for } \tilde{u} \right\}.$$

Now it is easy to see that the mapping  $f$  described above can be extended to an embedding of  $H$  into  $G$  if and only if the corresponding candidate graph has a matching covering  $\tilde{U}$ . Clearly, if the candidate graph  $B_f(H, G)$  of  $f$  has vertices  $\tilde{u} \in \tilde{U}$  of degree 0, then  $B_f(H, G)$  has no such matching and hence  $f$  cannot be extended. More generally we would like to avoid that  $\deg_{B_f(H, G)}(\tilde{u})$  is too small. Notice that this means precisely that  $f$  should not map  $N_H(\tilde{u})$  to a set  $B \subseteq V$  that has a small common neighbourhood in  $G$ . These sets  $B$  are the “bad” spots (see the beginning of this section) that should be avoided by  $f$ .

We explained above that, in order to avoid “bad” spots, we will have to change certain mappings  $f$  slightly. The exact definition of this operation is as follows.

**Definition 25** (switching). *Let  $f, f': X \rightarrow Y$  be injective functions. We say that  $f'$  is obtained from  $f$  by a switching if there are  $u, v \in X$  with  $f'(u) = f(v)$  and  $f'(v) = f(u)$  and  $f(w) = f'(w)$  for all  $w \notin \{u, v\}$ . The switching distance  $d_{\text{sw}}(f, f')$  of  $f$  and  $f'$  is at most  $s$  if the mapping  $f'$  can be obtained from  $f$  by a sequence of at most  $s$  switchings.*

These switchings will alter the candidate graph corresponding to the injective function slightly (but not much, see Lemma 27). In order to quantify this, we further define the neighbourhood distance between two bipartite graphs  $B$  and  $B'$  which determines the number of vertices (in one partition class) whose neighbourhoods differ in  $B$  and  $B'$ .

**Definition 26** (neighbourhood distance). *Let  $B = (U \dot{\cup} \tilde{U}, E)$ ,  $B' = (U \dot{\cup} \tilde{U}, E')$  be bipartite graphs. We define the neighbourhood distance of  $B$  and  $B'$  with respect to  $\tilde{U}$  as*

$$d_{N(\tilde{U})}(B, B') := |\{\tilde{u} \in \tilde{U} : N_B(\tilde{u}) \neq N_{B'}(\tilde{u})\}|.$$

The next simple lemma now examines the effect of switchings on the neighbourhood distance of candidate graphs and shows that functions with small switching distance correspond to candidate graphs with small neighbourhood distance.

**Lemma 27** (switching lemma). *Let  $H$  and  $G$  be bipartite graphs on vertex sets  $\tilde{U} \dot{\cup} \tilde{V}$  and  $U \dot{\cup} V$ , respectively, such that  $\deg_H(\tilde{v}) \leq \Delta$  for all  $\tilde{v} \in \tilde{V}$  and let  $f, f': \tilde{V} \rightarrow V$  be injective functions with switching distance  $d_{\text{sw}}(f, f') \leq s$ . Then the neighbourhood distance of the candidate graphs  $B_f(H, G)$  and  $B_{f'}(H, G)$  satisfies*

$$d_{N(\tilde{U})}(B_f(H, G), B_{f'}(H, G)) \leq 2s\Delta.$$

*Proof.* We proceed by induction on  $s$ . For  $s = 0$  the lemma is trivially true. Thus, consider  $s > 0$  and let  $g$  be a function with  $d_{\text{sw}}(f, g) \leq s - 1$  and  $d_{\text{sw}}(g, f') = 1$ . Define

$$N(f, f') := \left\{ \tilde{u} \in \tilde{U} : N_{B_f(H, G)}(\tilde{u}) \neq N_{B_{f'}(H, G)}(\tilde{u}) \right\}.$$

Clearly,  $|N(f, f')| = d_{N(\tilde{U})}(B_f(H, G), B_{f'}(H, G))$  and  $N(f, f') \subseteq N(f, g) \cup N(g, f')$ . By induction hypothesis we have  $|N(f, g)| \leq 2(s - 1)\Delta$ . The remaining switching from  $g$  to  $f'$  interchanges only the images of two vertices from  $\tilde{V}$ , say  $\tilde{v}_1$  and  $\tilde{v}_2$ . It follows that

$$N(g, f') = \left\{ \tilde{u} \in N_H(\tilde{v}_1) \cup N_H(\tilde{v}_2) : N_{B_g(H, G)}(\tilde{u}) \neq N_{B_{f'}(H, G)}(\tilde{u}) \right\},$$

which implies  $|N(g, f')| \leq 2\Delta$  and therefore we get  $|N(f, f')| \leq 2s\Delta$ .  $\square$

**11.2. A hypergraph packing lemma.** The main ingredient to the proof of the constrained blow-up lemma is the following hypergraph packing result (Lemma 29). To understand what this lemma says and how we will apply it, recall that we would like to embed the vertex set  $\tilde{U}$  of  $H$  into the vertex set  $U$  of  $G$  such that subsets of  $\tilde{U}$  that form neighbourhoods in the graph  $H$  avoiding certain “bad” spots in  $U$ . If  $H$  is a  $\Delta$ -regular graph, then these neighbourhoods form  $\Delta$ -sets. In this case, as we will see, also the “bad” spots form  $\Delta$ -sets. Accordingly, we have to solve the problem of packing the neighbourhood  $\Delta$ -sets  $\mathcal{N}$  and the “bad”  $\Delta$ -sets  $\mathcal{B}$ , which is a hypergraph packing problem. Lemma 29 below states that this is possible under certain conditions. One of these conditions is that the “bad” sets should not “cluster” too much (although there might be many of them). The following definition makes this precise.



**Definition 28** (corrupted sets). For  $\Delta \in \mathbb{N}$  and a set  $V$  let  $\mathcal{B} \subseteq \binom{V}{\Delta}$  be a collection of  $\Delta$ -sets in  $V$  and let  $x$  be a positive real. We say that all  $B \in \mathcal{B}$  are  $x$ -corrupted by  $\mathcal{B}$ . Recursively, for  $i \in [\Delta - 1]$  an  $i$ -set  $B \in \binom{V}{i}$  in  $V$  is called  $x$ -corrupted by  $\mathcal{B}$  if it is contained in more than  $x$  of the  $(i + 1)$ -sets that are  $x$ -corrupted by  $\mathcal{B}$ .

Observe that, if a vertex  $v \in V$  is not  $x$ -corrupted by  $\mathcal{B}$ , then it is also not  $x'$ -corrupted by  $\mathcal{B}$  for any  $x' > x$ .

The hypergraph packing lemma now implies that  $\mathcal{N}$  and  $\mathcal{B}$  can be packed if  $\mathcal{B}$  contains no corrupted sets. In fact this lemma states that *half of all* possible ways to map the vertices of  $\mathcal{N}$  to  $\mathcal{B}$  can be turned into such a packing by performing a sequence of few switchings.

**Lemma 29** (hypergraph packing lemma [25]). For all integers  $\Delta \geq 2$  and  $\ell \geq 1$  there are positive constants  $\eta_{29}$ , and  $n_{29}$  such that the following holds. Let  $\mathcal{B}$  be a  $\Delta$ -uniform hypergraph on  $n' \geq n_{29}$  vertices such that no vertex of  $\mathcal{B}$  is  $\eta_{29}n'$ -corrupted by  $\mathcal{B}$ . Let  $\mathcal{N}$  be a  $\Delta$ -uniform hypergraph on  $n \leq n'$  vertices such that no vertex of  $\mathcal{N}$  is contained in more than  $\ell$  edges of  $\mathcal{N}$ .

Then for at least half of all injective functions  $f: V(\mathcal{N}) \rightarrow V(\mathcal{B})$  there are packings  $f'$  of  $\mathcal{N}$  and  $\mathcal{B}$  with switching distance  $d_{\text{sw}}(f, f') \leq \sigma n$ .  $\square$

When applying this lemma we further make use of the following lemma which helps us to bound corruption.

**Lemma 30** (corruption lemma). Let  $n, \Delta > 0$  be integers and  $\mu$  and  $\eta$  be positive reals. Let  $V$  be a set of size  $n$  and  $\mathcal{B} \subseteq \binom{V}{\Delta}$  be a family of  $\Delta$ -sets of size at most  $\mu n^\Delta$ . Then at most  $(\Delta!/\eta^{\Delta-1})\mu n$  vertices are  $\eta n$ -corrupted by  $\mathcal{B}$ .

*Proof.* For  $i \in [\Delta]$  let  $\mathcal{B}_i$  be the family of all those  $i$ -sets  $B' \in \binom{V}{i}$  that are  $\eta n$ -corrupted by  $\mathcal{B}$ . We will prove by induction on  $i$  (starting at  $i = \Delta$ ) that

$$|\mathcal{B}_i| \leq \frac{\Delta!/i!}{\eta^{\Delta-i}} \mu n^i. \quad (32)$$

For  $i = 1$  this establishes the lemma. For  $i = \Delta$  the assertion is true by assumption. Now assume that (32) is true for  $i > 1$ . By definition every  $B' \in \mathcal{B}_{i-1}$  is contained in more than  $\eta n$  sets  $B \in \mathcal{B}_i$ . On the other hand, clearly every  $B \in \mathcal{B}_i$  contains at most  $i$  sets from  $\mathcal{B}_{i-1}$ . Double counting thus gives

$$\eta n |\mathcal{B}_{i-1}| \leq |\{(B', B) : B' \in \mathcal{B}_{i-1}, B \in \mathcal{B}_i, B' \subseteq B\}| \leq i |\mathcal{B}_i| \stackrel{(32)}{\leq} i \frac{\Delta!/i!}{\eta^{\Delta-i}} \mu n^i,$$

which implies (32) for  $i$  replaced by  $i - 1$ .  $\square$

**11.3. A matching lemma.** We indicated earlier that we are interested in determining whether a candidate graph has a matching covering one of its partition classes. In order to do so we will make use of the following matching lemma which is an easy consequence of Hall's theorem. This lemma takes two graphs  $B$  and  $B'$  as input that have small neighbourhood distance. In our application these two graphs will be candidate graphs that correspond to two injective mappings  $f$  and  $f'$  with small switching distance (such as promised by the hypergraph packing lemma, Lemma 29). Recall that Lemma 27 guarantees that mappings with small switching distance correspond to candidate graphs with small neighbourhood distance.

The matching lemma asserts that  $B'$  has the desired matching if certain vertex degree and neighbourhood conditions are satisfied. These conditions are somewhat technical. They are tailored exactly to match the conditions that we establish for candidate graphs in the proof of the constrained blow-up lemma (see Claims 34–36).

**Lemma 31** (matching lemma). Let  $B = (\tilde{U} \dot{\cup} U, E)$  and  $B' = (\tilde{U} \dot{\cup} U, E')$  be bipartite graphs with  $|U| \geq |\tilde{U}|$  and  $d_{N(\tilde{U})}(B, B') \leq s$ . If there are positive integers  $x$  and  $n_1, n_2, n_3$  such that

- (i)  $\deg_{B'}(\tilde{u}) \geq n_1$  for all  $\tilde{u} \in \tilde{U}$ ,
- (ii)  $|N_{B'}(\tilde{S})| \geq x|\tilde{S}|$  for all  $\tilde{S} \subseteq \tilde{U}$  with  $|\tilde{S}| \leq n_2$
- (iii)  $e_{B'}(\tilde{S}, S) \leq \frac{n_1}{n_3} |\tilde{S}| |S|$  for all  $\tilde{S} \subseteq \tilde{U}$ ,  $S \subseteq U$  with  $xn_2 \leq |S| < |\tilde{S}| < n_3$ ,

(iv)  $|N_B(S) \cap \tilde{S}| > s$  for all  $\tilde{S} \subseteq \tilde{U}$ ,  $S \subseteq U$  with  $|\tilde{S}| \geq n_3$  and  $|S| > |U| - |\tilde{S}|$ , then  $B'$  has a matching covering  $\tilde{U}$ .

*Proof.* We will check Hall's condition in  $B'$  for all sets  $\tilde{S} \subseteq \tilde{U}$ . We clearly have  $|N_{B'}(\tilde{S})| \geq |\tilde{S}|$  for  $|\tilde{S}| \leq xn_2$  by (ii) (if  $|\tilde{S}| > n_2$ , then consider a subset of  $\tilde{S}$  of size  $n_2$ ).

Next, consider the case  $xn_2 < |\tilde{S}| < n_3$ . Set  $S := N_{B'}(\tilde{S})$  and assume, for a contradiction, that  $|S| < |\tilde{S}|$ . Since  $|S| < |\tilde{S}| < n_3$  we have  $|S|/n_3 < 1$ . Therefore, applying (i), we can conclude that

$$e_{B'}(\tilde{S}, S) = \sum_{\tilde{u} \in \tilde{S}} |N_{B'}(\tilde{u})| \geq n_1 |\tilde{S}| > \frac{n_1}{n_3} |\tilde{S}| |S|,$$

which is a contradiction to (ii). Thus  $|N_{B'}(\tilde{S})| \geq |\tilde{S}|$ .

Finally, for sets  $\tilde{S}$  of size at least  $n_3$  set  $S := U \setminus N_{B'}(\tilde{S})$  and assume, again for a contradiction, that  $|N_{B'}(\tilde{S})| < |\tilde{S}|$ . This implies  $|S| > |U| - |\tilde{S}|$ . Accordingly we can apply (iv) to  $\tilde{S}$  and  $S$  and infer that  $|N_B(S) \cap \tilde{S}| > s$ . Since  $d_{N(\tilde{U})}(B, B') \leq s$ , at most  $s$  vertices from  $\tilde{U}$  have different neighbourhoods in  $B$  and  $B'$  and so

$$\begin{aligned} |N_{B'}(S) \cap \tilde{S}| &= \left| \left\{ \tilde{u} \in \tilde{S} : N_{B'}(\tilde{u}) \cap S \neq \emptyset \right\} \right| \\ &\geq \left| \left\{ \tilde{u} \in \tilde{S} : N_B(\tilde{u}) \cap S \neq \emptyset \right\} \right| - s = |N_B(S) \cap \tilde{S}| - s > 0, \end{aligned}$$

which is a contradiction as  $S = U \setminus N_{B'}(\tilde{S})$ .  $\square$

**11.4. Proof of Lemma 12.** Now we are almost ready to present the proof of the constrained blow-up lemma (Lemma 12). We just need one further technical lemma as preparation. This lemma considers a family of pairwise disjoint  $\Delta$ -sets  $\mathcal{S}$  in a set  $S$  and states that a random injective function from  $S$  to a set  $T$  usually has the following property. The images  $f(\mathcal{S})$  of sets in  $\mathcal{S}$  “almost” avoid a small family of “bad” sets  $\mathcal{T}$  in  $T$ .

**Lemma 32.** *For all positive integers  $\Delta$  and positive reals  $\beta$  and  $\mu_S$  there is  $\mu_T > 0$  such that the following holds. Let  $S$  and  $T$  be disjoint sets,  $\mathcal{S} \subseteq \binom{S}{\Delta}$  be a family of pairwise disjoint  $\Delta$ -sets in  $S$  with  $|\mathcal{S}| \leq \frac{1}{\Delta}(1 - \mu_S)|T|$ , and  $\mathcal{T} \subseteq \binom{T}{\Delta}$  be a family of  $\Delta$ -sets in  $T$  with  $|\mathcal{T}| \leq \mu_T |T|^\Delta$ .*

*Then a random injective function  $f: S \rightarrow T$  satisfies  $|f(\mathcal{S}) \setminus \mathcal{T}| > (1 - \beta)|\mathcal{S}|$  with probability at least  $1 - \beta^{|\mathcal{S}|}$ .*

*Proof.* Given  $\Delta$ ,  $\beta$ , and  $\mu_S$  choose

$$\mu_T := \sqrt[\beta]{\beta} \left( \frac{e}{\beta} \left( \frac{\Delta}{\mu_S} \right)^\Delta \right)^{-1}. \quad (33)$$

Let  $S$ ,  $T$ ,  $\mathcal{S}$ , and  $\mathcal{T}$  be as required and let  $f$  be a random injective function from  $S$  to  $T$ . We consider  $f$  as a consecutive random selection (without replacement) of images for the elements of  $S$  where the images of the elements of the (disjoint) sets in  $\mathcal{S}$  are chosen first. Let  $S_i$  be the  $i$ -th such set in  $\mathcal{S}$ . Then the probability that  $f$  maps  $S_i$  to a set in  $\mathcal{T}$ , which we denote by  $p_i$ , is at most

$$p_i \leq \frac{|\mathcal{T}|}{\binom{|T| - (i-1)\Delta}{\Delta}} \leq \frac{\mu_T |T|^\Delta}{\binom{\mu_S |T|}{\Delta}} \leq \mu_T \frac{|T|^\Delta}{\left( \frac{\mu_S |T|}{\Delta} \right)^\Delta} = \mu_T \left( \frac{\Delta}{\mu_S} \right)^\Delta =: p,$$

where the second inequality follows from  $(i-1)\Delta \leq |\bigcup \mathcal{S}| \leq (1 - \mu_S)|T|$ . Let  $Z$  be a random variable with distribution  $\text{Bi}(|\mathcal{S}|, p)$ . It follows that  $\mathbb{P}[|f(\mathcal{S}) \cap \mathcal{T}| \geq z] \leq \mathbb{P}[Z \geq z]$ . Since

$$\mathbb{P}[Z \geq z] \leq \binom{|\mathcal{S}|}{z} p^z < \left( \frac{e|\mathcal{S}|p}{z} \right)^z,$$

we infer that

$$\mathbb{P}[|f(\mathcal{S}) \cap \mathcal{T}| \geq \beta|\mathcal{S}|] < \left( \frac{ep}{\beta} \right)^{\beta|\mathcal{S}|} = \left( \frac{e\mu_T}{\beta} \left( \frac{\Delta}{\mu_S} \right)^\Delta \right)^{\beta|\mathcal{S}|} \stackrel{(33)}{=} \beta^{|\mathcal{S}|},$$

which proves the lemma since  $|f(\mathcal{S}) \cap \mathcal{T}| \geq \beta|\mathcal{S}|$  holds iff  $|f(\mathcal{S}) \setminus \mathcal{T}| \leq (1 - \beta)|\mathcal{S}|$ .  $\square$

Now we can finally give the proof of Lemma 12.

*Proof of Lemma 12.* We first define a sequence of constants. Given  $\Delta$ ,  $d$ , and  $\eta$  fix  $\Delta' := \Delta^2 + 1$ . Choose  $\beta$  and  $\sigma$  such that

$$\beta^{\frac{1}{7}(\frac{d}{2})^\Delta} \leq \frac{1}{5} \quad \text{and} \quad \frac{(1 - \beta)d^\Delta}{100^\Delta} \geq 2\sigma \quad (34)$$

Apply the hypergraph packing lemma, Lemma 29, with input  $\Delta$ ,  $\ell = 2\Delta + 1$ , and  $\sigma$  to obtain constants  $\eta_{29}$ , and  $n_{29}$ . Next, choose  $\eta'_{29}$ ,  $\mu_{\text{BL}}$ , and  $\mu_S$  such that

$$\frac{\eta'_{29}}{1 - \eta} \leq \eta_{29}, \quad \frac{\Delta! \cdot 2\mu_{\text{BL}}}{(\eta'_{29})^{\Delta-1}} \leq \eta, \quad \frac{1}{\Delta'} \leq \frac{1}{\Delta}(1 - \mu_S). \quad (35)$$

Lemma 32 with input  $\Delta$ ,  $\beta$ ,  $\mu_S$  provides us with a constant  $\mu_T$ . We apply Lemma 18 two times, once with input  $\Delta = \ell$ ,  $d$ ,  $\varepsilon' := \frac{1}{2}d$ , and  $\mu = \mu_{\text{BL}}/\Delta'$  and once with input  $\Delta = \ell$ ,  $d$ ,  $\varepsilon' := \frac{1}{2}d$ , and  $\mu = \mu_T$  and get constants  $\varepsilon_{18}$  and  $\tilde{\varepsilon}_{18}$ , respectively. Now we can fix the promised constant  $\varepsilon$  such that

$$\varepsilon \leq \min \left\{ \frac{\varepsilon_{18}}{\Delta'}, \frac{d}{2\Delta} \right\}, \quad \text{and} \quad \frac{\varepsilon\Delta'}{\eta(1 - \eta)} < \min\{d, \tilde{\varepsilon}_{18}\}. \quad (36)$$

As last input let  $r_1$  be given and set

$$\xi_{18} := \eta(1 - \eta)/(r_1\Delta'). \quad (37)$$

Next let  $c_{18}$  be the maximum of the two constants obtained from the two applications of Lemma 18, that we started above, with the additional parameter  $\xi_{18}$ . Further, let  $\nu$  and  $c_{19}$  be the constants from Lemma 19 for input  $\Delta$ ,  $d$ , and  $\varepsilon$ , and let  $c_{15}$  be the constant from Lemma 15 for input  $\Delta$  and  $\nu$ . Finally, we choose  $c = \max\{c_{18}, c_{19}, c_{15}\}$ . With this we defined all necessary constants.

Now assume we are given any  $1 \leq r \leq r_1$ , and a random graph  $\Gamma = \mathcal{G}_{n,p}$  with  $p \geq c(\log n/n)^{1/\Delta}$ , where, without loss of generality,  $n$  is such that

$$(1 - \eta')^{\frac{n}{r}} \geq n_{29}. \quad (38)$$

Then, with high probability, the graph  $\Gamma$  satisfies the assertion of the different lemmas concerning random graphs, that we started to apply in the definition of the constants. More precisely, by the choice of the constants above,

- (P1)  $\Gamma$  satisfies the assertion of Lemma 15 for parameters  $\Delta$  and  $\nu$ , i.e., for any set  $X$  and any family  $\mathcal{F}$  with the conditions required in this lemma, the conclusion of the lemma holds.
- (P2) Similarly  $\Gamma$  satisfies the assertion of Lemma 18 for parameters  $\Delta = \ell$ ,  $d$ ,  $\varepsilon' = \frac{1}{2}d$ ,  $\mu = \mu_{\text{BL}}/\Delta'$ ,  $\varepsilon_{18}$ , and  $\xi_{18}$ . The same holds for parameters  $\Delta = \ell$ ,  $d$ ,  $\varepsilon' = \frac{1}{2}d$ ,  $\mu = \mu_T$ ,  $\tilde{\varepsilon}_{18}$ , and  $\xi_{18}$ .
- (P3)  $\Gamma$  satisfies the assertion of Lemma 19 for parameters  $\Delta$ ,  $d$ ,  $\varepsilon$ , and  $\nu$ .

In the following we will assume that  $\Gamma$  has these properties and show that it then also satisfies the conclusion of the constrained blow-up lemma, Lemma 12.

Let  $G \subseteq \Gamma$  and  $H$  be two bipartite graphs on vertex sets  $U \dot{\cup} V$  and  $\tilde{U} \dot{\cup} \tilde{V}$ , respectively, that fulfil the requirements of Lemma 12. Moreover, let  $\mathcal{H} \subseteq \binom{\tilde{V}}{\Delta}$  be the family of special  $\Delta$ -sets, and  $\mathcal{B} \subseteq \binom{\tilde{V}}{\Delta}$  be the family of forbidden  $\Delta$ -sets. It is not difficult to see that, by possibly adding some edges to  $H$ , we can assume that the following holds.

- ( $\tilde{U}$ ) All vertices in  $\tilde{U}$  have degree *exactly*  $\Delta$ .
- ( $\tilde{V}$ ) All vertices in  $\tilde{V}$  have degree maximal  $\Delta + 1$ .

Our next step will be to split the partition class  $U$  of  $G$  and the corresponding partition class  $\tilde{U}$  of  $H$  into  $\Delta'$  parts of equal size. From the partition of  $H$  we require that no two vertices in one part have a common neighbour. This will guarantee that the neighbourhoods of two different vertices from one part form disjoint vertex sets (which we need because we would like to apply Lemma 19 later, in the proof of Claim 34, and Lemma 19 asserts certain properties for families of disjoint vertex sets).

Let us now explain precisely how we split  $U$  and  $\tilde{U}$ . We assume for simplicity that  $|\tilde{U}|$  and  $|U|$  are divisible by  $\Delta'$  and partition the sets  $U$  arbitrarily into  $\Delta'$  parts  $U = U_1 \dot{\cup} \dots \dot{\cup} U_{\Delta'}$  of equal

size, i.e., sets of size at least  $n/(r\Delta')$ . Similarly let  $\tilde{U} = \tilde{U}_1 \dot{\cup} \dots \dot{\cup} \tilde{U}_{\Delta'}$  be a partition of  $\tilde{U}$  into sets of equal size such that each  $\tilde{U}_j$  is 2-independent in  $H$ . Such a partition exists by the Theorem of Hajnal and Szemerédi (Theorem 22) applied to  $H^2[\tilde{U}]$  because the maximum degree of  $H^2$  is less than  $\Delta' = \Delta^2 + 1$ .

In Claim 33 below we will assert that there is an embedding  $f'$  of  $\tilde{V}$  into  $V$  that can be extended to *each* of the  $\tilde{U}_j$  separately such that we obtain an embedding of  $H$  into  $G$ . To this end we will consider the candidate graphs  $B_{f'}(H_j, G_j)$  defined by  $f'$  (see Definition 24) and show, that there is an  $f'$  such that each  $B_{f'}(H_j, G_j)$  has a matching covering  $\tilde{U}_j$ . This, as discussed earlier, will ensure the existence of the desired embedding. For preparing this argument, we first need to exclude some vertices of  $V$  which are not suitable for such an embedding. For identifying these vertices, we define the following family of  $\Delta$ -sets which contains  $\mathcal{B}$  and all sets in  $V$  that have a small common neighbourhood in some  $\tilde{U}_j$ .

Define  $\mathcal{B}' := \mathcal{B} \cup \bigcup_{j \in \Delta'} \mathcal{B}_j$  where

$$\mathcal{B}_j := \left\{ B \in \binom{V}{\Delta} : |N_G^\cap(B) \cap U_j| < (\tfrac{1}{2}d)^\Delta p^\Delta |U_j| \right\} \stackrel{(4)}{=} \text{bad}_{d/2, d, p}^{G, \Delta}(V, U_j). \quad (39)$$

We claim that we obtain a set  $\mathcal{B}'$  that is not much larger than  $\mathcal{B}$ . Indeed, by Proposition 6 the pair

$$(V, U_j) \text{ is } (\varepsilon\Delta', d, p)\text{-dense for all } j \in [\Delta'], \quad (40)$$

and  $\varepsilon\Delta' \leq \varepsilon_{18}$  by (36). Moreover we have  $|U_j| \geq n/(r\Delta') \geq n/(r_1\Delta') \geq \xi_{18}n$  by (37). We can thus use the fact that our random graph  $\Gamma$  satisfies property (P2) (with  $\mu = \mu_{\text{BL}}/\Delta'$ ) on the bipartite subgraph  $G[V \dot{\cup} U_j]$  and conclude that  $|\mathcal{B}_j| \leq \mu_{\text{BL}}|V|^\Delta/\Delta'$ . Since  $|\mathcal{B}| \leq \mu_{\text{BL}}|V|^\Delta$  by assumption we infer

$$|\mathcal{B}'| \leq \mu_{\text{BL}}|V|^\Delta + \Delta' \cdot \mu_{\text{BL}}|V|^\Delta/\Delta' = 2\mu_{\text{BL}}|V|^\Delta.$$

Set

$$V' := V \setminus V'' \quad \text{with} \quad V'' := \left\{ v \in V : v \text{ is } \eta'_{29}|V|\text{-corrupted by } \mathcal{B}' \right\} \quad (41)$$

and delete all sets from  $\mathcal{B}'$  that contain vertices from  $V''$ . This determines the set  $V''$  of vertices that we exclude from  $V$  for the embedding. We will next show that we did not exclude too many vertices in this process. For this we use the corruption lemma, Lemma 30. Indeed, Lemma 30 applied with  $n$  replaced by  $|V|$ , with  $\Delta, \mu = 2\mu_{\text{BL}}$ , and  $\eta'_{29}$  to  $V$  and  $\mathcal{B}'$  implies that

$$|V''| \leq \frac{\Delta!}{(\eta'_{29})^{\Delta-1}} 2\mu_{\text{BL}}|V| \stackrel{(35)}{\leq} \eta|V| \quad \text{and thus} \quad n' := |V'| \geq (1 - \eta)|V|. \quad (42)$$

Let

$$H_j := H[\tilde{U}_j \dot{\cup} \tilde{V}] \quad \text{and} \quad G_j := G[U_j \dot{\cup} V'].$$

Now we are ready to state the claim announced above, which asserts that there is an embedding  $f'$  of the vertices in  $\tilde{V}$  to the vertices in  $V'$  such that the corresponding candidate graphs  $B_{f'}(H_j, G_j)$  have matchings covering  $\tilde{U}_j$ . As we will shall show, this claim implies the assertion of the constrained blow-up lemma. Its proof, which we will provide thereafter, requires the matching lemma (Lemma 31), and the hypergraph packing lemma (Lemma 29).

**Claim 33.** *There is an injection  $f' : \tilde{V} \rightarrow V'$  with  $f'(T) \notin \mathcal{B}$  for all  $T \in \mathcal{H}$  such that for all  $j \in [\Delta']$  the candidate graph  $B_{f'}(H_j, G_j)$  has a matching covering  $\tilde{U}_j$ .*

Let us show that proving this claim suffices to establish the constrained blow-up lemma. Indeed, let  $f' : \tilde{V} \rightarrow V'$  be such an injection and denote by  $M_j : \tilde{U}_j \rightarrow U_j$  the corresponding matching in  $B_{f'}(H_j, G_j)$  for  $j \in [\Delta]$ . We claim that the function  $g : \tilde{U} \dot{\cup} \tilde{V} \rightarrow U \dot{\cup} V$ , defined by

$$g(\tilde{w}) = \begin{cases} M_j(\tilde{w}) & \tilde{w} \in \tilde{U}_j, \\ f'(\tilde{w}) & \tilde{w} \in \tilde{V}, \end{cases}$$

is an embedding of  $H$  into  $G$ . To see this, notice first that  $g$  is injective since  $f'$  is an injection and all  $M_j$  are matchings. Furthermore, consider an edge  $\tilde{u}\tilde{v}$  of  $H$  with  $\tilde{u} \in \tilde{U}_j$  for some  $j \in [\Delta']$  and  $\tilde{v} \in \tilde{V}$  and let

$$u := g(\tilde{u}) = M_j(\tilde{u}) \quad \text{and} \quad v := g(\tilde{v}) = f'(\tilde{v}).$$

It follows from the definition of  $M_j$  that  $\tilde{u}u$  is an edge of the candidate graph  $B_{f'}(H_j, G_j)$ . Hence, by the definition of  $B_{f'}(H_j, G_j)$ ,  $u$  is an  $f'$ -candidate for  $\tilde{u}$ , i.e.,

$$f'(N_{H_j}(\tilde{u})) \subseteq N_{G_j}(u).$$

Since  $v = f'(\tilde{v}) \in f'(N_{H_j}(\tilde{u}))$  this implies that  $uv$  is an edge of  $G$ . Because  $f'$  also satisfies  $f'(T) \notin \mathcal{B}$  for all  $T \in \mathcal{H}$  the embedding  $g$  also meets the remaining requirement of the constrained blow-up lemma that no special  $\Delta$ -set is mapped to a forbidden  $\Delta$ -set.  $\square$

For completing the proof of Lemma 12, we still need to prove Claim 33 which we shall be occupied with for the remainder of this section. We will assume throughout that we have the same setup as in the preceding proof. In particular all constants, sets, and graphs are defined as there.

For proving Claim 33 we will use the matching lemma (Lemma 31) on candidate graphs  $B = B_f(H_j, G_j)$  and  $B' = B_{f'}(H_j, G_j)$  for injections  $f, f': \tilde{V} \rightarrow V'$ . As we will see, the following three claims imply that there are suitable  $f$  and  $f'$  such that the conditions of this lemma are satisfied. More precisely, Claim 34 will take care of conditions (i) and (ii) in this lemma, Claim 35 of condition (iii), and Claim 36 of condition (iv). Before proving these claims we will show that they imply Claim 33.

The first claim states that *many* injective mappings  $f: \tilde{V} \rightarrow V'$  can be turned into injective mappings  $f'$  (with the help of a few switchings) such that the candidate graphs  $B_{f'}(H_j, G_j)$  for  $f'$  satisfy certain degree and expansion properties.

**Claim 34.** *For at least half of all injections  $f: \tilde{V} \rightarrow V'$  there is an injection  $f': \tilde{V} \rightarrow V'$  with  $d_{\text{sw}}(f, f') \leq \sigma n/r$  such that the following is satisfied for all  $j \in [\Delta']$ . For all  $\tilde{u} \in \tilde{U}_j$  and all  $\tilde{S} \subseteq \tilde{U}_j$  with  $|\tilde{S}| \leq p^{-\Delta}$  we have*

$$\deg_{B_{f'}(H_j, G_j)}(\tilde{u}) \geq \left(\frac{d}{2}\right)^\Delta p^\Delta |U_j| \quad \text{and} \quad |N_{B_{f'}(H_j, G_j)}(\tilde{S})| \geq \nu n p^\Delta |\tilde{S}|. \quad (43)$$

*Further, no special  $\Delta$ -set from  $\mathcal{H}$  is mapped to a forbidden  $\Delta$ -set from  $\mathcal{B}$  by  $f'$ .*

The second claim asserts that *all* injective mappings  $f'$  are such that the candidate graphs  $B_{f'}(H_j, G_j)$  do not contain sets of certain sizes with too many edges between them.

**Claim 35.** *All injections  $f': \tilde{V} \rightarrow V'$  satisfy the following for all  $j \in [\Delta']$  and all  $S \subseteq U_j$ ,  $\tilde{S} \subseteq \tilde{U}_j$ . If  $\nu n \leq |S| < |\tilde{S}| < \frac{1}{7} \left(\frac{d}{2}\right)^\Delta |U_j|$ , then*

$$e_{B_{f'}(H_j, G_j)}(\tilde{S}, S) \leq 7p^\Delta |\tilde{S}| |S|.$$

The last of the three claims states that for *random* injective mappings  $f$  the graphs  $B_{f'}(H_j, G_j)$  have edges between any pair of large enough sets  $S \subseteq U_j$  and  $\tilde{S} \subseteq \tilde{U}_j$ .

**Claim 36.** *A random injection  $f: \tilde{V} \rightarrow V'$  a.a.s. satisfies the following. For all  $j \in [\Delta']$  and all  $S \subseteq U_j$ ,  $\tilde{S} \subseteq \tilde{U}_j$  with  $|\tilde{S}| \geq \frac{1}{7} \left(\frac{d}{2}\right)^\Delta |U_j|$  and  $|S| > |U_j| - |\tilde{S}|$  we have*

$$|N_{B_f(H_j, G_j)}(S) \cap \tilde{S}| > 2\sigma n/r.$$

*Proof of Claim 33.* Our aim is to apply the matching lemma (Lemma 31) to the candidate graphs  $B_f(H_j, G_j)$  and  $B_{f'}(H_j, G_j)$  for all  $j \in [\Delta']$  with carefully chosen injections  $f$  and  $f'$ .

Let  $f: \tilde{V} \rightarrow V'$  be an injection satisfying the assertions of Claim 34 and Claim 36 and let  $f'$  be the injection promised by Claim 34 for this  $f$ . Such an  $f$  exists as at least half of all injections from  $\tilde{V}$  to  $V'$  satisfy the assertion of Claim 34 and almost all of those satisfy the assertion of

Claim 36. We will now show that for all  $j \in [\Delta']$  the conditions of Lemma 31 are satisfied for input

$$\begin{aligned} B &= B_f(H_j, G_j), & B' &= B_{f'}(H_j, G_j), & s &= 2\sigma n/r, \\ x &= \nu n p^\Delta, & n_1 &= (\tfrac{d}{2})^\Delta p^\Delta |U_j|, & n_2 &= p^{-\Delta}, & n_3 &= \tfrac{1}{7}(\tfrac{d}{2})^\Delta |U_j|, \end{aligned}$$

Claim 34 asserts that  $d_{\text{sw}}(f, f') \leq \sigma n/r$ . Since  $\tilde{U}_j$  is 2-independent in  $H$  we have  $\deg_{H_j}(\tilde{v}) \leq 1$  for all  $\tilde{v} \in \tilde{V}$ . Thus the switching lemma, Lemma 27, applied to  $H_j$  and  $G_j$  and with  $s$  replaced by  $\sigma n/r$  implies

$$d_{N(\tilde{U}_j)}(B, B') = d_{N(\tilde{U}_j)}(B_f(H_j, G_j), B_{f'}(H_j, G_j)) \leq 2\sigma n/r = s$$

Moreover, by Claim 34, for all  $\tilde{u} \in \tilde{U}_j$  we have

$$\deg_{B'}(\tilde{u}) = \deg_{B_{f'}(H_j, G_j)}(\tilde{u}) \geq (\tfrac{d}{2})^\Delta p^\Delta |U_j| = n_1$$

and thus condition (i) of Lemma 31 holds true. Further, we conclude from Claim 34 that  $|N_{B'}(\tilde{S})| \geq x|\tilde{S}|$  for all  $\tilde{S} \subseteq \tilde{U}_j$  with  $|\tilde{S}| < p^{-\Delta} = n_2$ . This gives condition (ii) of Lemma 31. In addition, Claim 35 states that for all  $S \subseteq U_j$ ,  $\tilde{S} \subseteq \tilde{U}_j$  with  $xn_2 = \nu n \leq |S| < |\tilde{S}| < \tfrac{1}{7}(\tfrac{d}{2})^\Delta |U_j| = n_3$  we have

$$e_{B'}(\tilde{S}, S) = e_{B_{f'}(H_j, G_j)}(\tilde{S}, S) \leq 7p^\Delta |\tilde{S}| |S| = \frac{n_1}{n_3} |\tilde{S}| |S|$$

and accordingly also condition (iii) of Lemma 31 is satisfied. To see (iv), observe that the choice of  $f$  and Claim 36 assert

$$|N_B(S) \cap \tilde{S}| = |N_{B_f(H_j, G_j)}(S) \cap \tilde{S}| > 2\sigma n/r = s$$

for all  $S \subseteq U_j$ ,  $\tilde{S} \subseteq \tilde{U}_j$  with  $|\tilde{S}| \geq \tfrac{1}{7}(\tfrac{d}{2})^\Delta |U_j| = n_3$  and  $|S| > |U| - |\tilde{S}|$ . Therefore, all conditions of Lemma 31 are satisfied and we infer that for all  $j \in [\Delta']$  the candidate graph  $B_{f'}(H_j, G_j)$  with  $f'$  as chosen above has a matching covering  $\tilde{U}$ . Moreover, by Claim 34,  $f'$  maps no special  $\Delta$ -set to a forbidden  $\Delta$ -set. This establishes Claim 33.  $\square$

It remains to show Claims 34–36. We start with Claim 34. For the proof of this claim we apply the hypergraph packing lemma (Lemma 29).

*Proof of Claim 34.* Notice that  $(\tilde{U})$  on page 29 implies that  $N_H(\tilde{u})$  contains exactly  $\Delta$  elements for each  $\tilde{u} \in \tilde{U}$ . Hence we may define the following family of  $\Delta$ -sets. Let

$$\mathcal{N} := \left\{ N_H(\tilde{u}) : \tilde{u} \in \tilde{U} \right\} \cup \mathcal{H} \subseteq \binom{\tilde{V}}{\Delta}.$$

We want to apply the hypergraph packing lemma (Lemma 29) with  $\Delta$ , with  $\ell$  replaced by  $2\Delta + 1$ , and with  $\sigma$  to the hypergraphs with vertex sets  $\tilde{V}$  and  $V'$  and edge sets  $\mathcal{N}$  and  $\mathcal{B}'$ , respectively (see (39) on page 30). We will first check that the necessary conditions are satisfied.

Observe that

$$|V'| \stackrel{(42)}{\geq} (1 - \eta')|V| \geq (1 - \eta')n/r \stackrel{(38)}{\geq} n_{29}, \quad \text{and} \quad |\tilde{V}| \leq |V'|.$$

Furthermore, a vertex  $\tilde{v} \in \tilde{V}$  is neither contained in more than  $\Delta$  sets from  $\mathcal{H}$  nor is  $\tilde{v}$  in  $N_H(\tilde{u})$  for more than  $\Delta + 1$  vertices  $\tilde{u} \in \tilde{U}$  (by  $(\tilde{V})$  on page 29). Therefore the condition Lemma 29 imposes on  $\mathcal{N}$  is satisfied with  $\ell$  replaced by  $2\Delta + 1$ . Moreover, according to (41) no vertex in  $V'$  is  $\eta'_{29}|V|$ -corrupted by  $\mathcal{B}'$ . Since

$$\eta'_{29}|V| \stackrel{(42)}{\leq} \eta'_{29}(1 - \eta)^{-1} n' \stackrel{(35)}{\leq} \eta_{29} n',$$

this (together with the observation in Definition 28) implies that no vertex in  $V'$  is  $\eta_{29} n'$ -corrupted by  $\mathcal{B}'$  and therefore all prerequisites of Lemma 29 are satisfied.

It follows that the conclusion of Lemma 29 holds for at least half of all injective functions  $f: \tilde{V} \rightarrow V'$ , namely that there are packings  $f'$  of (the hypergraphs with edges)  $\mathcal{N}$  and  $\mathcal{B}$  with switching distance  $d_{\text{sw}}(f, f') \leq \sigma|\tilde{V}| \leq \sigma n/r$ . Clearly, such a packing  $f'$  does not send any special



$\Delta$ -set from  $\mathcal{H}$  to any forbidden  $\Delta$ -set from  $\mathcal{B}$ . Our next goal is to show that  $f'$  satisfies the first part of (43) for all  $j \in [\Delta']$  and  $\tilde{u} \in \tilde{U}_j$ . For this purpose, fix  $j$  and  $\tilde{u}$ . The definition of the candidate graph  $B_{f'}(H_j, G_j)$ , Definition 24, implies

$$\begin{aligned} \deg_{B_{f'}(H_j, G_j)}(\tilde{u}) &= \left| \left\{ u \in U_j : f'(N_{H_j}(\tilde{u})) \subseteq N_{G_j}(u) \right\} \right| \\ &= \left| \left\{ u \in U_j : u \in N_{G_j}^\cap(f'(N_{H_j}(\tilde{u}))) \right\} \right| \\ &= \left| N_{G_j}^\cap(f'(N_{H_j}(\tilde{u}))) \right| \geq (\tfrac{1}{2}d)^\Delta p^\Delta |U_j|. \end{aligned}$$

where the first inequality follows from the fact that  $N_{H_j}(\tilde{u}) \in \mathcal{N}$  and thus, as  $f'$  is a packing of  $\mathcal{N}$  and  $\mathcal{B}'$ , we have  $f'(N_{H_j}(\tilde{u})) \notin \text{bad}_{d/2, d, p}^{G, \Delta}(V, U_j) \subseteq \mathcal{B}'$  (see the definition of  $\mathcal{B}'$  in (39)). This in turn means that all  $\Delta$ -sets  $f'(N_{H_j}(\tilde{u}))$  with  $\tilde{u} \in \tilde{U}_j$  are  $p$ -good (see Definition 17) in  $(V, U_j)$ , because  $(V, U_j)$  has  $p$ -density at least  $d - \varepsilon \Delta' \geq \frac{d}{2}$  by (40) and (36). With this information at hand we can proceed to prove the second part of (43). Let  $\tilde{S} \subseteq \tilde{U}_j$  with  $|\tilde{S}| < 1/p^\Delta$  and consider the family  $\mathcal{F} \subseteq \binom{V}{\Delta}$  with

$$\mathcal{F} := \{f'(N_H(\tilde{u})) : \tilde{u} \in \tilde{S}\}.$$

Because  $U_j$  is 2-independent in  $H$  the sets  $N_H(\tilde{u})$  with  $\tilde{u} \in \tilde{S}$  form a family of disjoint  $\Delta$ -sets in  $\tilde{V}$ . It follows that also the sets  $f'(N_H(\tilde{u}))$  with  $\tilde{u} \in \tilde{S}$  form a family of disjoint  $\Delta$ -sets in  $V$ . By (P3) on page 29 the conclusion of Lemma 19 holds for  $\Gamma$ . We conclude that the pair  $(V, U_j)$  is  $(1/p^\Delta, \nu n p^\Delta)$ -expanding. Since  $|\mathcal{F}| = |\tilde{S}| < 1/p^\Delta$  by assumption and all members of  $\mathcal{F}$  are  $p$ -good in  $(V, U_j)$  this implies that  $|N_{U_j}^\cap(\mathcal{F})| \geq \nu n p^\Delta |\mathcal{F}|$ . On the other hand  $N_{U_j}^\cap(\mathcal{F}) = N_{B_{f'}(H_j, G_j)}(\tilde{S})$  by the definition of  $B_{f'}(H_j, G_j)$  and  $\mathcal{F}$  and thus we get the second part of (43).  $\square$

Recall that property (P1) states that  $\Gamma$  satisfies the conclusion of Lemma 15 for certain parameters. We will use this fact to prove Claim 35.

*Proof of Claim 35.* Fix  $f' : \tilde{V} \rightarrow V'$ ,  $j \in [\Delta']$ ,  $S \subseteq U_j$ , and  $\tilde{S} \subseteq \tilde{U}_j$  with  $\nu n \leq |S| < |\tilde{S}| < \frac{1}{7}(\frac{d}{2})^\Delta |U_j|$ . For the candidate graphs  $B_{f'}(H_j, G_j)$  of  $f'$  we have

$$\begin{aligned} e_{B_{f'}(H_j, G_j)}(\tilde{S}, S) &= \left| \left\{ \tilde{u}u \in \tilde{S} \times S : f'(N_H(\tilde{u})) \subseteq N_G(u) \right\} \right| \\ &\stackrel{(2)}{=} \# \text{stars}^G \left( S, \{f'(N_H(\tilde{u})) : \tilde{u} \in \tilde{S}\} \right) \\ &\leq \# \text{stars}^\Gamma \left( S, \{f'(N_H(\tilde{u})) : \tilde{u} \in \tilde{S}\} \right) = \# \text{stars}^\Gamma(S, \mathcal{F}'), \end{aligned}$$

where  $\mathcal{F}' := \{f'(N_H(\tilde{u})) : \tilde{u} \in \tilde{S}\}$ . As before the sets  $f'(N_H(\tilde{u}))$  with  $\tilde{u} \in \tilde{S}$  form a family of  $|\tilde{S}|$  disjoint  $\Delta$ -sets in  $V'$ . Since  $\nu n \leq |S| < |\tilde{S}| = |\mathcal{F}'| \leq n$  we can appeal to property (P1) (and hence Lemma 15) with the set  $X := S$  and the family  $\mathcal{F}'$  and infer that

$$e_{B_{f'}(H_j, G_j)}(\tilde{S}, S) \leq \# \text{stars}^\Gamma(S, \mathcal{F}') \leq 7p^\Delta |\mathcal{F}'| |S| = 7p^\Delta |\tilde{S}| |S|$$

as required.  $\square$

Finally, we prove Claim 36. For this proof we will use the fact that  $\Delta$ -sets in  $p$ -dense graphs have big common neighbourhoods (the conclusion of Lemma 18 holds by property (P2)) together with Lemma 32.

*Proof of Claim 36.* Let  $f$  be an injective function from  $\tilde{V}$  to  $V'$ . First, consider a fixed  $j \in [\Delta']$  and fixed sets  $S \subseteq U_j$ ,  $\tilde{S} \subseteq \tilde{U}_j$  with  $|\tilde{S}| \geq \frac{1}{7}(\frac{d}{2})^\Delta |U_j|$  and  $|S| > |U_j| - |\tilde{S}|$ . Define

$$\mathcal{S} := \{N_{H_j}(\tilde{u}) : \tilde{u} \in \tilde{S}\} \quad \text{and} \quad \mathcal{T} := \text{bad}_{d/2, d, p}^{G, \Delta}(V', S).$$

and observe that

$$\begin{aligned} \left| N_{B_f(H_j, G_j)}(S) \cap \tilde{S} \right| &= \left| \left\{ \tilde{u} \in \tilde{S} : \exists u \in S \text{ with } f(N_{H_j}(\tilde{u})) \subseteq N_{G_j}(u) \right\} \right| \\ &= \left| \left\{ \tilde{u} \in \tilde{S} : N_{G_j}^\cap(f(N_{H_j}(\tilde{u}))) \cap S \neq \emptyset \right\} \right| \\ &\geq \left| \left\{ \tilde{u} \in \tilde{S} : f(N_{H_j}(\tilde{u})) \notin \text{bad}_{d/2, d, p}^{G, \Delta}(V', S) \right\} \right| = |f(S) \setminus \mathcal{T}| \end{aligned}$$

since all  $\Delta$ -sets  $B \notin \text{bad}_{d/2, d, p}^{G, \Delta}(V', S)$  satisfy  $|N_{G_j}^\cap(B) \cap S| \geq (\frac{d}{2})^\Delta p^\Delta |S| > 0$ . Thus, for proving the claim, it suffices to show that a random injection  $f : \tilde{V} \rightarrow V'$  violates  $|f(S) \setminus \mathcal{T}| > 2\sigma n/r$  with probability at most  $5^{-|U_j|}$  because this implies that  $f$  violates the conclusion of Claim 36 for *some*  $j \in [\Delta']$ , and *some*  $S \subseteq U_j$ ,  $\tilde{S} \subseteq \tilde{U}_j$  with probability at most  $\mathcal{O}(2^{|U_j|} 2^{|\tilde{U}_j|} \cdot 5^{-|U_j|}) = o(1)$ . For this purpose, we will use the fact that the pair  $(V', S)$  is  $p$ -dense. Indeed, observe that

$$|S| > |U_j| - |\tilde{S}| > |U_j| - |\tilde{U}_j| = \frac{|U| - |\tilde{U}|}{\Delta'} \geq \frac{\eta|U|}{\Delta'}$$

by the assumptions of the constrained blow-up lemma, Lemma 12. As  $|V'| \geq (1-\eta)|V|$  by (42) we can apply Proposition 6 twice to infer from the  $(\varepsilon, d, p)$ -density of  $(V, U)$  that  $(V', S)$  is  $(\tilde{\varepsilon}, d, p)$ -dense with  $\tilde{\varepsilon} := \varepsilon \Delta' / (\eta(1-\eta))$ . Furthermore  $\tilde{\varepsilon} \leq \tilde{\varepsilon}_{18}$  by (36) and

$$|V'| \stackrel{(42)}{\geq} (1-\eta) \frac{n}{r} \stackrel{(37)}{\geq} \xi_{18} n, \quad \text{and} \quad |S| > \frac{\eta|U|}{\Delta'} \geq \frac{\eta n}{r \Delta'} \stackrel{(37)}{\geq} \xi_{18} n.$$

Hence we conclude from (P2) on page 29 (with  $\mu = \mu_T$ ) that  $|\mathcal{T}| = |\text{bad}_{d/2, d, p}^{G, \Delta}(V', S)| \leq \mu_T |V'|^\Delta$ . In addition

$$\frac{1}{7} \left(\frac{d}{2}\right)^\Delta |U_j| \leq |\tilde{S}| = |S| \leq |\tilde{U}_j| \leq (1-\eta) \frac{n}{\Delta'} \stackrel{(42)}{\leq} \frac{|V'|}{\Delta'} \stackrel{(35)}{\leq} \frac{1}{\Delta} (1-\mu_S) |V'|. \quad (44)$$

Thus, we can apply Lemma 32 with  $\Delta$ ,  $\beta$ , and  $\mu_S$  to  $S = \tilde{V}$ ,  $T = V'$ , and to  $\mathcal{S}$  and  $\mathcal{T}$  and conclude that  $f$  violates

$$|f(\mathcal{S}) \setminus \mathcal{T}| > (1-\beta) |S| \stackrel{(44)}{\geq} (1-\beta) \frac{1}{7} \left(\frac{d}{2}\right)^\Delta |U_j| \geq \frac{(1-\beta) d^\Delta n}{7 \cdot 2^{\Delta} r \Delta'} \geq \frac{(1-\beta) d^\Delta n}{100 \Delta r} \stackrel{(34)}{\geq} 2\sigma \frac{n}{r}$$

with probability at most

$$\beta^{|S|} \leq \beta^{\frac{1}{7} \left(\frac{d}{2}\right)^\Delta |U_j|} \leq 5^{-|U_j|}$$

where the first inequality follows from (44) and the second from (34).  $\square$

## REFERENCES

- [1] M. Albert, A. Frieze, and B. Reed, *Multicoloured Hamilton cycles*, Electron. J. Combin. **2** (1995), 13 pp.
- [2] N. Alon and M. Capalbo, *Sparse universal graphs for bounded-degree graphs*, Random Structures Algorithms **31** (2007), 123–133.
- [3] ———, *Optimal universal graphs with deterministic embedding*, Proc. of the 19th Annual ACM-SIAM Symp. on Discrete Algorithms SODA, 2008, pp. 373–378.
- [4] N. Alon, M. Capalbo, Y. Kohayakawa, V. Rödl, A. Ruciński, and E. Szemerédi, *Universality and tolerance*, Proc. of the 41st Annual Symp. on Foundations of Computer Science FOCS, 2000, pp. 14–21.
- [5] N. Alon, M. Krivelevich, and B. Sudakov, *Embedding nearly-spanning bounded degree trees*, Combinatorica **27** (2007), no. 6, 629–644.
- [6] N. Alon and Z. Füredi, *Spanning subgraphs of random graphs*, Graphs Combin. **8** (1992), no. 1, 91–94.
- [7] J. Balogh, B. Csaba, and W. Samotij, *Local resilience of almost spanning trees in random graphs*, Submitted.
- [8] J. Böttcher, K. P. Pruessmann, A. Taraz, and A. Würfl, *Bandwidth, expansion, treewidth, separators, and universality for bounded degree graphs*, European J. Combin., Accepted for publication.
- [9] J. Böttcher, M. Schacht, and A. Taraz, *Proof of the bandwidth conjecture of Bollobás and Komlós*, Math. Ann. **343** (2009), no. 1, 175–205.
- [10] D. Dellamonica, Y. Kohayakawa, M. Marcinišzyn, and A. Steger, *On the resilience of long cycles in random graphs*, Electron. J. Combin. **15** (2008), 26 pp., R32.
- [11] D. Dellamonica, Y. Kohayakawa, V. Rödl, and A. Ruciński, *Universality of random graphs*, Proc. of the 19th Annual ACM-SIAM Symp. on Discrete Algorithms SODA, 2008, pp. 782–788.
- [12] G. A. Dirac, *Some theorems on abstract graphs*, Proc. London Math. Soc. (3) **2** (1952), 69–81.
- [13] P. Erdős, J. Nešetřil, and V. Rödl, *On some problems related to partitions of edges of a graph*, Graphs and other combinatorial topics (Prague, 1982), vol. 59, Teubner, Leipzig, 1983, pp. 54–63.

- [14] A. Frieze and B. Reed, *Polychromatic Hamilton cycles*, Discrete Math. **118** (1993), no. 1-3, 69–74.
- [15] S. Gerke, Y. Kohayakawa, V. Rödl, and A. Steger, *Small subsets inherit sparse  $\epsilon$ -regularity*, J. Combin. Theory Ser. B **97** (2007), no. 1, 34–56.
- [16] S. Gerke and A. Steger, *The sparse regularity lemma and its applications*, Surveys in combinatorics 2005, London Math. Soc. Lecture Note Ser., vol. 327, Cambridge Univ. Press, Cambridge, 2005, pp. 227–258.
- [17] G. Hahn and C. Thomassen, *Path and cycle sub-Ramsey numbers and an edge-colouring conjecture*, Discrete Math. **62** (1986), no. 1, 29–33.
- [18] A. Hajnal and E. Szemerédi, *Proof of a conjecture of P. Erdős*, Combinatorial theory and its applications, II (Proc. Colloq., Balatonfüred, 1969), North-Holland, Amsterdam, 1970, pp. 601–623.
- [19] S. Janson, T. Luczak, and A. Ruciński, *Random graphs*, Wiley-Interscience, New York, 2000.
- [20] Y. Kohayakawa, *Szemerédi’s regularity lemma for sparse graphs*, Foundations of computational mathematics (Rio de Janeiro, 1997), Springer, 1997, pp. 216–230.
- [21] Y. Kohayakawa and V. Rödl, *Regular pairs in sparse random graphs. I*, Random Structures Algorithms **22** (2003), no. 4, 359–434.
- [22] ———, *Szemerédi’s regularity lemma and quasi-randomness*, Recent advances in algorithms and combinatorics, CMS Books Math./Ouvrages Math. SMC., vol. 11, Springer, 2003, pp. 289–35.
- [23] Y. Kohayakawa, M. Schacht, V. Rödl, and E. Szemerédi, *Sparse partition universal graphs for graphs of bounded degree*, submitted, 2008.
- [24] V. Rödl and A. Ruciński, *Perfect matchings in  $\epsilon$ -regular graphs and the blow-up lemma*, Combinatorica **19** (1999), no. 3, 437–452.
- [25] V. Rödl, A. Ruciński, and A. Taraz, *Hypergraph packing and graph embedding*, Combinatorics, Probability and Computing **8** (1999), 363–376.
- [26] B. Sudakov and V. Vu, *Local resilience of graphs*, Random Structures Algorithms **33** (2008), 409–433.
- [27] P. Turán, *Eine Extremalaufgabe aus der Graphentheorie*, Mat. Fiz. Lapok **48** (1941), 436–452.

#### APPENDIX A. THE CONNECTION LEMMA

The proof of Lemma 13 which we present in this appendix is inherent in the proof of [23, Lemma 18]. The only difference is that we have a somewhat more special set-up here (given by the pre-defined partitions and candidate sets). This set-up however is chosen exactly in such a way that this proof continues to work if we adapt the parameters involved accordingly.

*Proof of Lemma 13.* For the proof of Lemma 13 we use an inductive argument and embed a partition class of  $H$  into the corresponding partition class of  $G$  one at a time. Before describing this strategy we will define two graph properties  $D_p(d_0, \epsilon', \mu, \epsilon, \xi)$  and  $\text{STAR}_p(k, \xi, \nu)$ , which a random graph  $\Gamma = \mathcal{G}_{n,p}$  enjoys a.a.s. for suitable sets of parameters. Then we will set up these parameters accordingly and define all other constants involved in the proof.

For a fixed  $n$ -vertex graph  $\Gamma$ , fixed positive reals  $d_0, \epsilon', \mu, \epsilon, \xi$ , and  $\nu$ , a fixed integer  $k$ , and a function  $p = p(n)$  we define the following properties of  $\Gamma$ .

$D_p(d_0, \epsilon', \mu, \epsilon, \xi)$ : We say that  $\Gamma$  has property  $D_p(d_0, \epsilon', \mu, \epsilon, \xi)$  if it satisfies the property stated in Lemma 20 with these parameters and with  $\Delta$ , i.e., whenever  $G = (X \dot{\cup} Y \dot{\cup} Z, E)$  is a tripartite subgraph of  $\Gamma$  with the required properties, then it satisfies the conclusion of this lemma.

$\text{STAR}_p(k, \xi, \nu)$ : Similarly  $\Gamma$  has property  $\text{STAR}_p(k, \xi, \nu)$  if  $\Gamma$  has the property stated in Lemma 16 with  $\Delta$  replaced by  $k$ , with parameters  $\xi, \nu$ , and for  $p = p(n)$ .

Now we set up the constants. Let  $\Delta, t$  and  $d$  be given and assume without loss of generality that  $d \leq \frac{1}{4}$ . First we set

$$\mu = \frac{1}{4\Delta^2} \tag{45}$$

and we fix  $\epsilon_i$  for  $i = t, t-1, \dots, 0$  by setting

$$\begin{aligned} \epsilon_t &= \frac{d}{12\Delta t}, \quad d_0 := d, \quad \text{and} \\ \epsilon_{i-1} &= \min \{ \epsilon(\Delta, d_0, \epsilon' = \epsilon_i, \mu), \epsilon_i \} \text{ for } i = t, \dots, 1, \end{aligned} \tag{46}$$

where  $\epsilon(\Delta - 1, d_0, \epsilon' = \epsilon_i, \mu)$  is given by Lemma 20. We choose  $\epsilon := \epsilon_0$  and  $\xi := (d/100)^\Delta$  and receive  $r_1$  as input. For each  $k \in [\Delta]$  and each  $r' \in [r_1]$  Lemma 16 with  $\Delta$  replaced by  $k$  and with  $\xi$  replaced by  $\xi/r'$  provides positive constants  $\nu(k, r')$  and  $c(k, r')$ . Let  $\nu$  be the minimum among the  $\nu(k, r')$  and let  $c_{16}$  be the maximum among the  $c(k, r')$  as we let both  $k$  and  $r'$  vary. Similarly Lemma 20 with input  $\Delta - 1, d_0, \epsilon' = \epsilon_i, \mu$ , and  $\xi$  replaced by  $\xi/r'$  provides constants

$c'(i, r')$  for  $i \in [0, t]$  and  $r' \in [r_1]$ . We let  $c_{20}$  be the maximum among these  $c'(i, r')$ . Then we fix  $c := \max\{c_{16}, c_{20}\}$ , and receive  $r \in [r_1]$  as input. Finally, we set

$$\xi_{16} := \xi_{20} := \xi/r = (d/100)^\Delta (1/r). \quad (47)$$

This finishes the definition of the constants.

Let  $p = p(n) \geq c(\log n/n)^{1/\Delta}$  and let  $\Gamma$  be a graph from  $\mathcal{G}_{n,p}$ . By Lemma 16, Lemma 20, and the choice of constants the graph  $\Gamma$  a.a.s. satisfies properties  $D_p(d, \varepsilon_i, \mu, \varepsilon_{i-1}, \xi_{20})$  for all  $i \in [t]$ , and properties  $\text{STAR}_p(k, \xi_{16}, \nu)$  for all  $k \in [\Delta]$ . In the remainder of this proof we assume that  $\Gamma$  has these properties and show that then  $\Gamma$  also satisfies the conclusion of Lemma 13.

Let  $G \subseteq \Gamma$  and  $H$  be arbitrary graphs satisfying the requirements stated in the lemma on vertex sets  $W = W_1 \dot{\cup} \dots \dot{\cup} W_t$  and  $\widetilde{W} = \widetilde{W}_1 \dot{\cup} \dots \dot{\cup} \widetilde{W}_t$ , respectively. Let  $h: \widetilde{W} \rightarrow [t]$  be the “partition function” for the vertex partition of  $H$ , i.e.,

$$h(\widetilde{w}) = j \quad \text{if and only if} \quad \widetilde{w} \in \widetilde{W}_j.$$

For an integer  $i \leq h(\widetilde{w})$  we denote by

$$\text{ldeg}^i(\widetilde{w}) := |N_H(\widetilde{w}) \cap \{\widetilde{x} \in \widetilde{W} : h(\widetilde{x}) \leq i\}|$$

the left degree of  $\widetilde{w}$  with respect to  $\widetilde{W}_1 \dot{\cup} \dots \dot{\cup} \widetilde{W}_i$ . Clearly  $\text{ldeg}^{h(\widetilde{w})}(\widetilde{w}) = \text{ldeg}(\widetilde{w})$ . Before we continue, recall that each vertex  $\widetilde{w} \in \widetilde{W}_i$  is equipped with a set  $X_{\widetilde{w}} \subseteq V(\Gamma) \setminus W$  and that we defined an external degree  $\text{edeg}(\widetilde{w}) = |X_{\widetilde{w}}|$  of  $\widetilde{w}$  as well as a candidate set  $C(\widetilde{w}) = N_{\widetilde{W}_i}^\cap(X_{\widetilde{w}}) \subseteq W_i$ . In the course of our embedding procedure, that we will describe below, we shall shrink this candidate set but keep certain invariants as we explain next.

We proceed inductively and embed the vertex class  $\widetilde{W}_i$  into  $W_i$  one at a time, for  $i = 1, \dots, t$ . To this end, we verify the following statement  $(\mathcal{S}_i)$  for  $i = 0, \dots, t$ .

- $(\mathcal{S}_i)$  There exists a partial embedding  $\varphi_i$  of  $H[\bigcup_{j=1}^i \widetilde{W}_j]$  into  $G[\bigcup_{j=1}^i W_j]$  such that for every  $\widetilde{z} \in \bigcup_{j=i+1}^t \widetilde{W}_j$  there exists a candidate set  $C_i(\widetilde{z}) \subseteq C(\widetilde{z})$  given by
- (a)  $C_i(\widetilde{z}) = \bigcap \{N_G(\varphi_i(\widetilde{x})) : \widetilde{x} \in N_H(\widetilde{z}) \text{ and } h(\widetilde{x}) \leq i\} \cap C(\widetilde{z})$ ,  
and satisfying
  - (b)  $|C_i(\widetilde{z})| \geq (dp/2)^{\text{ldeg}^i(\widetilde{z})} |C(\widetilde{z})|$ , and
  - (c) for every edge  $\{\widetilde{z}, \widetilde{z}'\} \in E(H)$  with  $h(\widetilde{z}), h(\widetilde{z}') > i$  the pair  $(C_i(\widetilde{z}), C_i(\widetilde{z}'))$  is  $(\varepsilon_i, d, p)$ -dense in  $G$ .

Statement  $(\mathcal{S}_i)$  ensures the existence of a partial embedding of the first  $i$  classes  $\widetilde{W}_1, \dots, \widetilde{W}_i$  of  $H$  into  $G$  such that for every unembedded vertex  $\widetilde{z}$  there exists a candidate set  $C_i(\widetilde{z}) \subseteq C(\widetilde{z})$  that is not too small (see part (b)). Moreover, if we embed  $\widetilde{z}$  into its candidate set, then its image will be adjacent to all vertices  $\varphi_i(\widetilde{x})$  with  $\widetilde{x} \in (\widetilde{W}_1 \dot{\cup} \dots \dot{\cup} \widetilde{W}_i) \cap N_H(\widetilde{z})$  (see part (a)). The last property, part (c), says that for edges of  $H$  such that none of the endvertices are embedded already the respective candidate sets induce  $(\varepsilon, d', p)$ -dense pairs for some positive  $d'$ . This property will be crucial for the inductive proof.

*Remark.* In what follows we shall use the following convention. Since the embedding of  $H$  into  $G$  will be divided into  $t$  rounds, we shall find it convenient to distinguish among the vertices of  $H$ . We shall use  $\widetilde{x}$  for vertices that have already been embedded,  $\widetilde{y}$  for vertices that will be embedded in the current round, while  $\widetilde{z}$  will denote vertices that we shall embed at a later step.

Before we verify  $(\mathcal{S}_i)$  for  $i = 0, \dots, t$  by induction on  $i$  we note that  $(\mathcal{S}_t)$  implies that  $H$  can be embedded into  $G$  in such a way that every vertex  $\widetilde{w} \in \widetilde{W}$  is mapped to a vertex in its candidate set  $C(\widetilde{w})$ . Consequently, verifying  $(\mathcal{S}_t)$  concludes the proof of Lemma 13.

Basis of the induction:  $i = 0$ . We first verify  $(\mathcal{S}_0)$ . In this case  $\varphi_0$  is the empty mapping and for every  $\widetilde{z} \in \widetilde{W}$  we have, according to (a),  $C_0(\widetilde{z}) = C(\widetilde{z})$ , as there is no vertex  $\widetilde{x} \in N_H(\widetilde{z})$  with  $h(\widetilde{x}) \leq 0$ . Property (b) holds because  $C_0(\widetilde{z}) = C(\widetilde{z})$  and  $\text{ldeg}^0(\widetilde{z}) = 0$  for every  $\widetilde{z} \in \widetilde{W}$ . Finally, property (c) follows from the property that  $(C(\widetilde{z}), C(\widetilde{z}'))$  is  $(\varepsilon_0, d, p)$ -dense by (E) of Lemma 13.

Induction step:  $i \rightarrow i + 1$ . For the inductive step, we suppose that  $i < t$  and assume that statement  $(\mathcal{S}_i)$  holds; we have to construct  $\varphi_{i+1}$  with the required properties. Our strategy is as follows. In the first step, we find for every  $\tilde{y} \in \widetilde{W}_{i+1}$  an appropriate subset  $C'(\tilde{y}) \subseteq C_i(\tilde{y})$  of its candidate set such that if  $\varphi_{i+1}(\tilde{y})$  is chosen from  $C'(\tilde{y})$ , then the new candidate set  $C_{i+1}(\tilde{z}) := C_i(\tilde{z}) \cap N_G(\varphi_{i+1}(\tilde{y}))$  of every “right-neighbour”  $\tilde{z}$  of  $\tilde{y}$  will not shrink too much and property (c) will continue to hold.

Note, however, that in general  $|C'(\tilde{y})| \leq |C_i(\tilde{y})| = o(n) \ll |\widetilde{W}_{i+1}|$  (if  $\text{ldeg}^i(\tilde{y}) \geq 1$ ) and, hence, we cannot “blindly” select  $\varphi_{i+1}(\tilde{y})$  from  $C'(\tilde{y})$ . Instead, in the second step, we shall verify Hall’s condition to find a system of distinct representatives for the indexed set system  $(C'(\tilde{y}) : \tilde{y} \in \widetilde{W}_{i+1})$  and we let  $\varphi_{i+1}(\tilde{y})$  be the representative of  $C'(\tilde{y})$ . (A similar idea was used in [6, 24].) We now give the details of those two steps.

*First step:* For the first step, fix  $\tilde{y} \in \widetilde{W}_{i+1}$  and set

$$N_H^{i+1}(\tilde{y}) := \{\tilde{z} \in N_H(\tilde{y}) : h(\tilde{z}) > i + 1\}.$$

A vertex  $v \in C_i(\tilde{y})$  will be “bad” (i.e., we shall not select  $v$  for  $C'(\tilde{y})$ ) if there exists a vertex  $\tilde{z} \in N_H^{i+1}(\tilde{y})$  for which  $N_G(v) \cap C_i(\tilde{z})$ , in view of (b) and (c) of  $(\mathcal{S}_{i+1})$ , cannot play the rôle of  $C_{i+1}(\tilde{z})$ .

We first prepare for (b) of  $(\mathcal{S}_{i+1})$ . Fix a vertex  $\tilde{z} \in N_H^{i+1}(\tilde{y})$ . Since  $(C_i(\tilde{y}), C_i(\tilde{z}))$  is an  $(\varepsilon_i, d, p)$ -dense pair by (c) of  $(\mathcal{S}_i)$ , Proposition 7 implies that there exist at most  $\varepsilon_i |C_i(\tilde{y})| \leq \varepsilon_t |C_i(\tilde{y})|$  vertices  $v$  in  $C_i(\tilde{y})$  such that

$$|N_G(v) \cap C_i(\tilde{z})| < (d - \varepsilon_t)p |C_i(\tilde{z})|.$$

Repeating the above for all  $\tilde{z} \in N_H^{i+1}(\tilde{y})$ , we infer from (a) and (b) of  $(\mathcal{S}_i)$ , that there are at most  $\Delta \varepsilon_t |C_i(\tilde{y})|$  vertices  $v \in C_i(\tilde{y})$  such that the following fails to be true for some  $\tilde{z} \in N_H^{i+1}(\tilde{y})$ :

$$|N_G(v) \cap C_i(\tilde{z})| \geq (d - \varepsilon_t)p |C_i(\tilde{z})|$$

$$\stackrel{(a),(b)}{\geq} (d - \varepsilon_t)p \left(\frac{dp}{2}\right)^{\text{ldeg}^i(\tilde{z})} |C(\tilde{z})| \stackrel{(46)}{\geq} \left(\frac{dp}{2}\right)^{\text{ldeg}^{i+1}(\tilde{z})} |C(\tilde{z})|. \quad (48)$$

For property (c) of  $(\mathcal{S}_{i+1})$ , we fix an edge  $e = \{\tilde{z}, \tilde{z}'\}$  with  $h(\tilde{z}), h(\tilde{z}') > i + 1$  and with at least one end vertex in  $N_H^{i+1}(\tilde{y})$ . There are at most  $\Delta(\Delta - 1) < \Delta^2$  such edges. Note that if both vertices  $\tilde{z}$  and  $\tilde{z}'$  are neighbours of  $\tilde{y}$ , i.e.,  $\tilde{z}, \tilde{z}' \in N_H^{i+1}(\tilde{y})$ , then

$$\max \{ \text{ldeg}^i(\tilde{y}) + \text{edeg}(\tilde{y}), \text{ldeg}^i(\tilde{z}) + \text{edeg}(\tilde{z}), \text{ldeg}^i(\tilde{z}') + \text{edeg}(\tilde{z}') \} \leq \Delta - 2,$$

by (C) of Lemma 13 and because all three vertices  $\tilde{y}$ ,  $\tilde{z}$ , and  $\tilde{z}'$  have at least two neighbours in  $\widetilde{W}_{i+1} \cup \dots \cup \widetilde{W}_t$ . From property (b) of  $(\mathcal{S}_i)$ , and (A) and (D) of Lemma 13 we infer for all  $\tilde{w} \in \{\tilde{y}, \tilde{z}, \tilde{z}'\}$  that

$$|C_i(\tilde{w})| \stackrel{(b)}{\geq} \left(\frac{dp}{2}\right)^{\text{ldeg}^i(\tilde{w})} |C(\tilde{w})| \stackrel{(A),(D)}{\geq} \left(\frac{dp}{2}\right)^{\text{ldeg}^i(\tilde{w}) + \text{edeg}(\tilde{w})} \frac{n}{r} \stackrel{(47)}{\geq} \xi_{20} p^{\Delta-2} n.$$

Furthermore,  $\Gamma$  has property  $D_p(d, \varepsilon_{i+1}, \mu, \varepsilon_i, \xi_{20})$  by assumption. This implies that there are at most  $\mu |C_i(\tilde{y})|$  vertices  $v \in C_i(\tilde{y})$  such that the pair  $(N_G(v) \cap C_i(\tilde{z}), N_G(v) \cap C_i(\tilde{z}'))$  fails to be  $(\varepsilon_{i+1}, d, p)$ -dense.

If, on the other hand, say, only  $\tilde{z} \in N_H^{i+1}(\tilde{y})$  and  $\tilde{z}' \notin N_H^{i+1}(\tilde{y})$ , then

$$\max \{ \text{ldeg}^i(\tilde{y}) + \text{edeg}(\tilde{y}), \text{ldeg}^i(\tilde{z}') + \text{edeg}(\tilde{z}') \} \leq \Delta - 1$$

$$\text{and } \text{ldeg}^i(\tilde{z}) + \text{edeg}(\tilde{z}) \leq \Delta - 2.$$

Consequently, similarly as above,

$$\min \{ |C_i(\tilde{y})|, |C_i(\tilde{z}')| \} \geq \xi_{20} p^{\Delta-1} n \quad \text{and} \quad |C_i(\tilde{z})| \geq \xi_{20} p^{\Delta-2} n$$

and we can appeal to the fact that  $\Gamma$  has property  $D_p(d, \varepsilon_{i+1}, \mu, \varepsilon_i, \xi_{20})$  to infer that there are at most  $\mu |C_i(\tilde{y})|$  vertices  $v \in C_i(\tilde{y})$  such that  $(N_G(v) \cap C_i(\tilde{z}), C_i(\tilde{z}'))$  fails to be  $(\varepsilon_{i+1}, d, p)$ -dense.

For a given  $v \in C_i(\tilde{y})$ , let  $\hat{C}_i(\tilde{z}) = C_i(\tilde{z}) \cap N_G(v)$  if  $\tilde{z} \in N_H^{i+1}(\tilde{y})$  and  $\hat{C}_i(\tilde{z}) = C_i(\tilde{z})$  if  $\tilde{z} \notin N_H^{i+1}(\tilde{y})$ , and define  $\hat{C}_i(\tilde{z}')$  analogously.

Summarizing the above we infer that there are at least

$$(1 - \Delta\varepsilon_t - \Delta^2\mu)|C_i(\tilde{y})| \quad (49)$$

vertices  $v \in C_i(\tilde{y})$  such that

- (b')  $|N_G(v) \cap C_i(\tilde{z})| \geq (dp/2)^{\text{ldeg}^{i+1}(\tilde{z})}|C(\tilde{z})|$  for every  $\tilde{z} \in N_H^{i+1}(\tilde{y})$  (see (48)) and
- (c')  $(\hat{C}_i(\tilde{z}), \hat{C}_i(\tilde{z}'))$  is  $(\varepsilon_{i+1}, d, p)$ -dense for all edges  $\{\tilde{z}, \tilde{z}'\}$  of  $H$  with  $h(\tilde{z}), h(\tilde{z}') > i+1$  and  $\{\tilde{z}, \tilde{z}'\} \cap N_H^{i+1}(\tilde{y}) \neq \emptyset$ .

Let  $C'(\tilde{y})$  be the set of those vertices  $v$  from  $C_i(\tilde{y})$  satisfying properties (b') and (c') above. Recall that  $\text{ldeg}^i(\tilde{y}) + \text{edeg}(\tilde{y}) = \text{ldeg}^i(\tilde{y}') + \text{edeg}(\tilde{y}')$  for all  $\tilde{y}, \tilde{y}' \in \widetilde{W}_{i+1}$  and set

$$k := \text{ldeg}^i(\tilde{y}) + \text{edeg}(\tilde{y}) \text{ for some } \tilde{y} \in \widetilde{W}_{i+1}. \quad (50)$$

Since  $\tilde{y} \in \widetilde{W}_{i+1}$  was arbitrary, we infer from property (b) of  $(\mathcal{S}_i)$ , properties (A) and (D) of Lemma 13, and the choices of  $\mu$  and  $\varepsilon_t$  that

$$\begin{aligned} |C'(\tilde{y})| &\stackrel{(49)}{\geq} (1 - \Delta\varepsilon_t - \Delta^2\mu)|C_i(\tilde{y})| \stackrel{(b)}{\geq} (1 - \Delta\varepsilon_t - \Delta^2\mu) \left(\frac{dp}{2}\right)^{\text{ldeg}^i(\tilde{y})} |C(\tilde{z})| \\ &\stackrel{(A),(D)}{\geq} (1 - \Delta\varepsilon_t - \Delta^2\mu) \left(\frac{dp}{2}\right)^k \frac{n}{r} \stackrel{(45),(46)}{\geq} \left(\frac{dp}{10}\right)^k \frac{n}{r}. \end{aligned} \quad (51)$$

*Second step:* We now turn to the aforementioned second part of the inductive step. Here we ensure the existence of a system of distinct representatives for the indexed set system

$$\mathcal{C}_{i+1} := \left( C'(\tilde{y}) : \tilde{y} \in \widetilde{W}_{i+1} \right).$$

We shall appeal to *Hall's condition* and show that for every subfamily  $\mathcal{C}' \subseteq \mathcal{C}_{i+1}$  we have

$$|\mathcal{C}'| \leq \left| \bigcup_{C' \in \mathcal{C}'} C' \right|. \quad (52)$$

Because of (51), assertion (52) holds for all families  $\mathcal{C}'$  with  $1 \leq |\mathcal{C}'| \leq (dp/10)^k n/r$ .

Thus, consider a family  $\mathcal{C}' \subseteq \mathcal{C}_i$  with  $|\mathcal{C}'| > (dp/10)^k n/r$ . For every  $\tilde{y} \in \widetilde{W}_{i+1}$  we have a set  $\tilde{K}(\tilde{y})$  of  $\text{ldeg}^i(\tilde{y})$  already embedded vertices of  $H$  such that  $\tilde{K}(\tilde{y}) = N_H(\tilde{y}) \setminus N_H^{i+1}(\tilde{y})$ . Let  $K'(\tilde{y}) := \varphi_i(\tilde{K}(\tilde{y}))$  be the image of  $\tilde{K}(\tilde{y})$  in  $G$  under  $\varphi_i$ . Recall that  $\tilde{y}$  is equipped with a set  $X_{\tilde{y}} \subseteq V(\Gamma) \setminus W$  of size  $\text{edeg}(\tilde{y})$  in Lemma 13. We have  $\text{ldeg}^i(\tilde{y}) + \text{edeg}(\tilde{y}) = k$  by (50). Hence, when we add the vertices of  $X_{\tilde{y}}$  to  $K'(\tilde{y})$  we obtain a set  $K(\tilde{y}) = \{u_1(\tilde{y}), \dots, u_k(\tilde{y})\}$  of  $k$  vertices in  $\Gamma$ . Note that for two distinct vertices  $\tilde{y}, \tilde{y}' \in \widetilde{W}_{i+1}$  the sets  $\tilde{K}(\tilde{y})$  and  $\tilde{K}(\tilde{y}')$  are disjoint. This follows from the fact that the distance in  $H$  between  $\tilde{y}$  and  $\tilde{y}'$  is at least four by the 3-independence of  $\widetilde{W}_{i+1}$  (cf. (B) of Lemma 13) and if  $\tilde{K}(\tilde{y}) \cap \tilde{K}(\tilde{y}') \neq \emptyset$ , then this distance would be at most two. In addition  $(X_{\tilde{y}} : \tilde{y} \in \widetilde{W}_{i+1})$  consists of pairwise disjoint sets by hypothesis. Consequently, the sets  $K(\tilde{y})$  and  $K(\tilde{y}')$  are disjoint as well and, therefore,

$$\mathcal{F} := \{K(\tilde{y}) : C'(\tilde{y}) \in \mathcal{C}'\} \subseteq \{K(\tilde{y}) : \tilde{y} \in \widetilde{W}_{i+1}\} \subseteq \binom{V(\Gamma)}{k}$$

is a family of  $|\mathcal{C}'|$  pairwise disjoint  $k$ -sets in  $V(\Gamma)$ . Moreover,  $C(\tilde{y}) = N_{W_i}^\cap(X_{\tilde{y}})$  by definition and so (a) of  $(\mathcal{S}_i)$  implies

$$C'(\tilde{y}) \subseteq C(\tilde{y}) \cap \bigcap_{v \in K'(\tilde{y})} N_\Gamma(v) = \bigcap_{v \in K(\tilde{y})} N_\Gamma(v).$$

Let

$$U = \bigcup_{C'(\tilde{y}) \in \mathcal{C}'} C'(\tilde{y}) \subseteq W_{i+1},$$



and suppose for a contradiction that

$$|U| < |\mathcal{C}'| = |\mathcal{F}|. \quad (53)$$

We now use the fact that  $\Gamma$  has property  $\text{STAR}_p(k, \xi_{16}, \nu)$  and apply it to  $U$  and  $\mathcal{F}$  (see Lemma 16). By assumption  $|U| < |\mathcal{F}| \leq \nu n p^k |\mathcal{F}|$ . We deduce that

$$\# \text{stars}^\Gamma(U, \mathcal{F}) \leq p^k |U| |\mathcal{F}| + 6\xi_{16} n p^k |\mathcal{F}|.$$

On the other hand, because of (51), we have

$$\# \text{stars}^\Gamma(U, \mathcal{F}) \geq \left(\frac{dp}{10}\right)^k \frac{n}{r} |\mathcal{F}|.$$

Combining the last two inequalities we infer from property (A) of Lemma 13 that

$$|U| \geq \left(\left(\frac{d}{10}\right)^k \frac{1}{r} - 6\xi_{16}\right) n \stackrel{(47)}{\geq} \xi_{16} n \stackrel{(47)}{=} \xi \frac{n}{r} \stackrel{(A)}{\geq} |\widetilde{W}_{i+1}| \geq |\mathcal{C}'|,$$

which contradicts (53). This contradiction shows that (53) does not hold, that is, Hall's condition (52) does hold. Hence, there exists a system of representatives for  $\mathcal{C}_{i+1}$ , i.e., an injective mapping  $\psi: \widetilde{W}_{i+1} \rightarrow \bigcup_{\widetilde{y} \in \widetilde{W}_{i+1}} C'(\widetilde{y})$  such that  $\psi(\widetilde{y}) \in C'(\widetilde{y})$  for every  $\widetilde{y} \in \widetilde{W}_{i+1}$ .

Finally, we extend  $\varphi_i$ . For that we set

$$\varphi_{i+1}(\widetilde{w}) = \begin{cases} \varphi_i(\widetilde{w}), & \text{if } \widetilde{w} \in \bigcup_{j=1}^i \widetilde{W}_j, \\ \psi(\widetilde{w}), & \text{if } \widetilde{w} \in \widetilde{W}_{i+1}. \end{cases}$$

Note that every  $\widetilde{z} \in \bigcup_{j=i+2}^t \widetilde{W}_j$  has at most one neighbour in  $\widetilde{W}_{i+1}$ , as otherwise there would be two vertices  $\widetilde{y}$  and  $\widetilde{y}' \in \widetilde{W}_{i+1}$  with distance at most 2 in  $H$ , which contradicts property (B) of Lemma 13. Consequently, for every  $\widetilde{z} \in \bigcup_{j=i+2}^t \widetilde{W}_j$  we have

$$C_{i+1}(\widetilde{z}) = \begin{cases} C_i(\widetilde{z}), & \text{if } N_H(\widetilde{z}) \cap \widetilde{W}_{i+1} = \emptyset, \\ C_i(\widetilde{z}) \cap N_G(\varphi_{i+1}(\widetilde{y})), & \text{if } N_H(\widetilde{z}) \cap \widetilde{W}_{i+1} = \{\widetilde{y}\}. \end{cases}$$

by (a) of  $(\mathcal{S}_{i+1})$ . In what follows we show that  $\varphi_{i+1}$  and  $C_{i+1}(\widetilde{z})$  for every  $\widetilde{z} \in \bigcup_{j=i+2}^t \widetilde{W}_j$  have the desired properties and validate  $(\mathcal{S}_{i+1})$ .

First of all, from (a) of  $(\mathcal{S}_i)$ , combined with  $\varphi_{i+1}(\widetilde{y}) \in C'(\widetilde{y}) \subseteq C_i(\widetilde{y})$  for every  $\widetilde{y} \in \widetilde{W}_{i+1}$  and the property that  $(\varphi_{i+1}(\widetilde{y}): \widetilde{y} \in \widetilde{W}_{i+1})$  is a system of distinct representatives, we infer that  $\varphi_{i+1}$  is indeed a partial embedding of  $H[\bigcup_{j=1}^{i+1} \widetilde{W}_j]$ .

Next we shall verify property (b) of  $(\mathcal{S}_{i+1})$ . So let  $\widetilde{z} \in \bigcup_{j=i+2}^t \widetilde{W}_j$  be fixed. If  $N_H(\widetilde{z}) \cap \widetilde{W}_{i+1} = \emptyset$ , then  $C_{i+1}(\widetilde{z}) = C_i(\widetilde{z})$ ,  $\text{ldeg}^{i+1}(\widetilde{z}) = \text{ldeg}^i(\widetilde{z})$ , which yields (b) of  $(\mathcal{S}_{i+1})$  for that case. If, on the other hand,  $N_H(\widetilde{z}) \cap \widetilde{W}_{i+1} \neq \emptyset$ , then there exists a unique neighbour  $\widetilde{y} \in \widetilde{W}_{i+1}$  of  $H$  (owing to the 3-independence of  $\widetilde{W}_{i+1}$  by property (B) of Lemma 13). As discussed above we have  $C_{i+1}(\widetilde{z}) = C_i(\widetilde{z}) \cap N_G(\varphi_{i+1}(\widetilde{y}))$  in this case. Since  $\varphi_{i+1}(\widetilde{y}) \in C'(\widetilde{y})$ , we infer directly from (b') that (b) of  $(\mathcal{S}_{i+1})$  is satisfied.

Finally, we verify property (c) of  $(\mathcal{S}_{i+1})$ . Let  $\{\widetilde{z}, \widetilde{z}'\}$  be an edge of  $H$  with  $\widetilde{z}, \widetilde{z}' \in \bigcup_{j=i+2}^t \widetilde{W}_j$ . We consider three cases, depending on the size of  $N_H(\widetilde{z}) \cap \widetilde{W}_{i+1}$  and of  $N_H(\widetilde{z}') \cap \widetilde{W}_{i+1}$ . If  $N_H(\widetilde{z}) \cap \widetilde{W}_{i+1} = \emptyset$  and  $N_H(\widetilde{z}') \cap \widetilde{W}_{i+1} = \emptyset$ , then part (c) of  $(\mathcal{S}_{i+1})$  follows directly from part (c) of  $(\mathcal{S}_i)$  and  $\varepsilon_{i+1} \geq \varepsilon_i$ , combined with  $C_{i+1}(\widetilde{z}) = C_i(\widetilde{z})$ ,  $C_{i+1}(\widetilde{z}') = C_i(\widetilde{z}')$ . If  $N_H(\widetilde{z}) \cap \widetilde{W}_{i+1} = \{\widetilde{y}\}$  and  $N_H(\widetilde{z}') \cap \widetilde{W}_{i+1} = \emptyset$ , then (c) of  $(\mathcal{S}_{i+1})$  follows from (c') and the definition of  $C_{i+1}(\widetilde{z})$  and  $C_{i+1}(\widetilde{z}')$ . If  $N_H(\widetilde{z}) \cap \widetilde{W}_{i+1} = \{\widetilde{y}\}$  and  $N_H(\widetilde{z}') \cap \widetilde{W}_{i+1} = \{\widetilde{y}'\}$ , then  $\widetilde{y} = \widetilde{y}'$ , as otherwise there would be a  $\widetilde{y}$ - $\widetilde{y}'$ -path in  $H$  with three edges, contradicting the 3-independence of  $\widetilde{W}_{i+1}$ . Consequently, (c) of  $(\mathcal{S}_{i+1})$  follows from (c') and the definition of  $C_{i+1}(\widetilde{z})$  and  $C_{i+1}(\widetilde{z}')$ .

We have therefore verified (a)–(c) of  $(\mathcal{S}_i)$ , thus concluding the induction step. The proof of Lemma 13 follows by induction.  $\square$

## APPENDIX B. PROOFS OF AUXILIARY LEMMAS

In this section we provide all proofs that were postponed earlier, namely those of Lemma 16, Lemma 18, and Lemma 21.

**B.1. Proof of Lemma 16.** This proof makes use of a Chernoff bound for the binomially distributed random variable  $\# \text{stars}^\Gamma(X, \mathcal{F})$  appearing in this lemma (cf. Definition 14 and the discussion below this definition).

*Proof of Lemma 16.* Given  $\Delta$  and  $\xi$  let  $\nu$  and  $c$  be constants satisfying

$$\begin{aligned} -6\xi \log(2\xi) &\leq -(6\xi - 2\sqrt{\nu}) \log \xi, & 2\nu &\leq (\sqrt{\nu} - 2\nu), \\ \Delta + 1 - 6\xi c^\Delta &\leq -1, & \text{and} & \Delta \leq \nu c^\Delta. \end{aligned} \quad (54)$$

First we estimate the probability that there are  $X$  and  $\mathcal{F}$  with  $|\mathcal{F}| \geq n/\log n$  fulfilling the requirements of the lemma but violating (3). Chernoff's inequality  $\mathbb{P}[Y \geq \mathbb{E}Y + t] \leq \exp(-t)$  for a binomially distributed random variable  $Y$  and  $t \geq 6\mathbb{E}Y$  (see [19, Chapter 2]): implies

$$\mathbb{P} \left[ \# \text{stars}^\Gamma(X, \mathcal{F}) \geq p^\Delta |X| |\mathcal{F}| + 6\xi np^\Delta |\mathcal{F}| \right] \leq \exp(-6\xi np^\Delta |\mathcal{F}|) \leq \exp(-6\xi c^\Delta |\mathcal{F}| \log n)$$

for fixed  $X$  and  $\mathcal{F}$  since  $6\xi np^\Delta |\mathcal{F}| \geq 6p^\Delta |X| |\mathcal{F}|$ . As the number of choices for  $\mathcal{F}$  and  $X$  can be bounded by  $\sum_{f=n/\log n}^{\xi n} n^{\Delta f}$  and  $2^n \leq \exp(n)$ , respectively, the probability we want to estimate is at most

$$\sum_{f=\frac{n}{\log n}}^{\xi n} \exp \left( \Delta f \log n + n - 6\xi c^\Delta f \log n \right) \leq \sum_{f=\frac{n}{\log n}}^{\xi n} \exp \left( f \log n (\Delta + 1 - 6\xi c^\Delta) \right),$$

which does not exceed  $\xi n \exp(-n)$  by (54) and thus tends to 0 as  $n$  tends to infinity.

It remains to establish a similar bound on the probability that there are  $X$  and  $\mathcal{F}$  with  $|\mathcal{F}| < n/\log n$  fulfilling the requirements of the lemma but violating (3). For this purpose we use that

$$\mathbb{P}[Y \geq t] \leq q^t \binom{m}{t} \leq \exp \left( -t \log \frac{t}{3qm} \right)$$

for a random variable  $Y$  with distribution  $\text{Bi}(m, q)$  and infer for fixed  $X$  and  $\mathcal{F}$

$$\begin{aligned} \mathbb{P} \left[ \# \text{stars}^\Gamma(X, \mathcal{F}) \geq p^\Delta |X| |\mathcal{F}| + 6\xi np^\Delta |\mathcal{F}| \right] &\leq \mathbb{P} \left[ \# \text{stars}^\Gamma(X, \mathcal{F}) \geq 6\xi np^\Delta |\mathcal{F}| \right] \\ &\leq \exp \left( -6\xi np^\Delta |\mathcal{F}| \log \frac{2\xi n}{|\mathcal{F}|} \right) \leq \exp \left( -2\sqrt{\nu} np^\Delta |\mathcal{F}| \log \frac{n}{|\mathcal{F}|} \right). \end{aligned}$$

because  $-6\xi \log(2\xi) \leq -(6\xi - 2\sqrt{\nu}) \log \xi \leq (6\xi - 2\sqrt{\nu}) \log(n/|X|)$  by (54). The number of choices for  $\mathcal{F}$  and  $X$  in total can be bounded by

$$\begin{aligned} \sum_{f=1}^{\frac{n}{\log n}} \sum_{x=1}^{\nu np^\Delta f} n^{\Delta f} \binom{n}{x} &\leq \sum_{f=1}^{\frac{n}{\log n}} \sum_{x=1}^{\nu np^\Delta f} \exp \left( \Delta f \log n + \nu np^\Delta f \log \frac{en}{x} \right) \\ &\leq \sum_{f=1}^{\frac{n}{\log n}} \sum_{x=1}^{\nu np^\Delta f} \exp \left( 2\nu np^\Delta f \log \frac{en}{x} \right) \leq \sum_{f=1}^{\frac{n}{\log n}} \sum_{x=1}^{\nu np^\Delta f} \exp \left( \sqrt{\nu} np^\Delta f \log \frac{n}{x} \right) \end{aligned}$$

where the second inequality follows from  $\Delta \log n \leq \nu c^\Delta \log n \leq \nu np^\Delta$  and the last from  $2\nu \log e \leq (\sqrt{\nu} - 2\nu) \log(n/x)$  by (54). Therefore the probability under consideration is at most

$$\sum_{f=1}^{\frac{n}{\log n}} \sum_{x=1}^{\nu np^\Delta f} \exp \left( \sqrt{\nu} np^\Delta f \log \frac{n}{x} - 2\sqrt{\nu} np^\Delta f \log \frac{n}{x} \right) \leq n^2 \exp \left( -\sqrt{\nu} \log n \frac{n}{\log n} \right).$$

□

**B.2. Proof of Lemma 18.** We will use the following simple proposition about cuts in hypergraphs. This proposition generalises the well known fact that any graph  $G$  admits a vertex partition into sets of roughly equal size such that the resulting cut contains at least half the edges of  $G$ .

**Proposition 37.** *Let  $\mathcal{G} = (V, \mathcal{E})$  be an  $\ell$ -uniform hypergraph with  $m$  edges and  $n$  vertices such that  $n \geq 3\ell$ . Then there is a partition  $V = V_1 \dot{\cup} V_2$  with  $|V_1| = \lfloor 2n/3 \rfloor$  and  $|V_2| = \lceil n/3 \rceil$  such that at least  $m \cdot \ell/2^{\ell+2}$  edges in  $\mathcal{E}$  are 1-crossing, i.e., they have exactly one vertex in  $V_2$ .*

*Proof.* Let  $X$  be the number of  $\frac{1}{3}$ -cuts of  $V$ , i.e., cuts  $V = V_1 \dot{\cup} V_2$  with  $|V_1| = \lfloor 2n/3 \rfloor$  and  $|V_2| = \lceil n/3 \rceil$ . For a fixed edge  $B$  there are precisely  $2^\ell$  ways to distribute its vertices over  $V_1 \dot{\cup} V_2$  out of which exactly  $\ell$  are such that  $B$  is 1-crossing. Further, for  $r$  fixed vertices of  $B$  exactly  $\binom{n-\ell}{\lceil n/3 \rceil - r}$  of all  $\frac{1}{3}$ -cuts have exactly these vertices in  $V_2$ . It is easy to check that

$$\binom{n-\ell}{\lceil n/3 \rceil - r} \leq 4 \binom{n-\ell}{\lceil n/3 \rceil - 1} \quad \text{for all } 0 \leq r \leq \ell.$$

It follows that  $B$  is 1-crossing for at least an  $\frac{1}{4}\ell/(2^\ell)$  fraction of all  $\frac{1}{3}$ -cuts. Now assume that all  $\frac{1}{3}$ -cuts have less than  $m \cdot \ell/2^{\ell+2}$  edges that are 1-crossing. Then double counting gives

$$m \cdot \frac{\ell}{2^{\ell+2}} \cdot X > \sum_{B \in \mathcal{E}} \#\left\{ \frac{1}{3}\text{-cuts s.t. } B \text{ is 1-crossing} \right\} \geq m \cdot \frac{1}{4} \frac{\ell}{2^\ell} \cdot X$$

which is a contradiction.  $\square$

In the proof of Lemma 18 we need to estimate the number of “bad”  $\ell$ -sets in a vertex set  $X$ . For this purpose we will use Proposition 37 to obtain a partition of  $X$  into sets  $X = X_1 \dot{\cup} X_2$  such that a substantial proportion of all these bad  $\ell$ -sets will be 1-crossing and  $X_1$  is not too small. In this way we obtain many  $(\ell - 1)$ -sets in  $X_1$  most of which will, as we show, be similarly bad as the  $\ell$ -sets we started with. This will allow us to prove Lemma 18 by induction.

*Proof of Lemma 18.* Let  $\Delta$  and  $d$  be given. Let  $\Gamma$  be an  $n$ -vertex graph, let  $\ell$  be an integer, let  $\varepsilon', \mu, \varepsilon, \xi$  be positive real numbers, and let  $p = p(n)$  be a function. We say that  $\Gamma$  has property  $P_\ell(\varepsilon', \mu, \varepsilon, \xi, p(n))$  if  $\Gamma$  has the property stated in Lemma 18 with parameters  $\varepsilon', \mu, \varepsilon, \xi, p(n)$  and with parameters  $\Delta$  and  $d$ . Similarly,  $\Gamma$  has property  $D(\varepsilon', \mu, \varepsilon, \xi, p(n))$  if it satisfies the conclusion of Lemma 20 with these parameters and with  $\Delta$  and  $d_0 := d$ . For any fixed  $\ell > 0$ , we denote by  $(\mathcal{P}_\ell)$  the following statement.

$(\mathcal{P}_\ell)$  *For all  $\varepsilon', \mu > 0$  there is  $\varepsilon$  such that for all  $\xi > 0$  there is  $c > 1$  such that a random graph  $\Gamma = \mathcal{G}_{n,p}$  with  $p > c(\frac{\log n}{n})^{1/\Delta}$  has property  $P_\ell(\varepsilon', \mu, \varepsilon, \xi, p(n))$  with probability  $1 - o(1)$ .*

We prove that  $(\mathcal{P}_\ell)$  holds for every fixed  $\ell > 0$  by induction on  $\ell$ . The case  $\ell = 1$  is an easy consequence of Proposition 7 which states that in all  $(\varepsilon, d, p)$ -dense pairs most vertices have a large neighbourhood.

For the inductive step assume that  $(\mathcal{P}_{\ell-1})$  holds. We will show that this implies  $(\mathcal{P}_\ell)$ . We start by specifying the constants appearing in statement  $(\mathcal{P}_\ell)$ . Let  $\varepsilon'$  and  $\mu$  be arbitrary positive constants. Set  $\varepsilon'_{\ell-1} := \varepsilon'$  and  $\mu_{\ell-1} := \frac{1}{100}\mu \frac{\ell}{2^{\ell+2}}$ . Let  $\varepsilon_{\ell-1}$  be given by  $(\mathcal{P}_{\ell-1})$  for input parameters  $\varepsilon'_{\ell-1}$  and  $\mu_{\ell-1}$ . Set  $\varepsilon'_{20} := \varepsilon_{\ell-1}$  and let  $\varepsilon_{20}$  be as promised by Lemma 20 with parameters  $\varepsilon'_{20}$  and  $\mu_{20} := \frac{1}{2}$ . Define  $\varepsilon := \mu_{\ell-1}\varepsilon_{20}\varepsilon'_{\ell-1}$ . Next, let  $\xi$  be an arbitrary parameter provided by the adversary in Lemma 18 and choose  $\xi_{\ell-1} := \xi(d - \varepsilon)$  and  $\xi_{20} := \mu_{\ell-1}\xi$ . Finally, let  $c_{\ell-1}$  and  $c_{20}$  be given by  $(\mathcal{P}_{\ell-1})$  and by Lemma 20, respectively, for the previously specified parameters together with  $\xi_{\ell-1}$  and  $\xi_{20}$ . Set  $c := \max\{c_{\ell-1}, c_{20}\}$ . We will prove that with this choice of  $\varepsilon$  and  $c$  the statement in  $(\mathcal{P}_\ell)$  holds for the input parameters  $\varepsilon', \mu$ , and  $\xi$ .

Let  $\Gamma = \mathcal{G}_{n,p}$  be a random graph. By  $(\mathcal{P}_{\ell-1})$  and Lemma 20, and by the choice of the parameters the graph  $\Gamma$  has properties

$$P_{\ell-1}(\varepsilon'_{\ell-1}, \mu_{\ell-1}, \varepsilon_{\ell-1}, \xi_{\ell-1}, p(n)) \quad \text{and} \quad D(\varepsilon'_{20}, \mu_{20}, \varepsilon_{20}, \xi_{20}, p(n))$$

with probability  $1 - o(1)$  if  $n$  is large enough. We will show that a graph  $\Gamma$  with these properties also satisfies  $P_\ell(\varepsilon', \mu, \varepsilon, \xi, p(n))$ . Let  $G = (X \dot{\cup} Y, E)$  be an arbitrary subgraph of such a  $\Gamma$  where  $|X| = n_1$  and  $|Y| = n_2$  with  $n_1 \geq \xi p^{\Delta-1}n$ ,  $n_2 \geq \xi p^{\Delta-\ell}n$ , and  $(X, Y)$  is an  $(\varepsilon, d, p)$ -dense pair.

We would like to show that for  $\mathcal{B}_\ell := \text{bad}_{\varepsilon', d, p}^{G, \ell}(X, Y)$  we have  $|\mathcal{B}_\ell| \leq \mu n_1^\ell$ . Assume for a contradiction that this is not the case. By Proposition 37 there is a cut  $X = X_1 \dot{\cup} X_2$  with  $|X_1| = \lfloor 2n_1/3 \rfloor$  and  $|X_2| = \lceil n_2/3 \rceil$  such that at least  $|\mathcal{B}_\ell| \cdot \ell/2^{\ell+2}$  of the  $\ell$ -sets in  $\mathcal{B}_\ell$  are 1-crossing, i.e., have exactly one vertex in  $X_2$ . By Proposition 7 there are less than  $\varepsilon|X|$  vertices  $x \in X_2$  such that  $|N_Y(x)| < (d - \varepsilon)pn_2$ . We delete all  $\ell$ -sets from  $\mathcal{B}_\ell$  that contain such a vertex or are not 1-crossing for  $X = X_1 \dot{\cup} X_2$  and call the resulting set  $\mathcal{B}'_\ell$ . It follows that

$$|\mathcal{B}'_\ell| \geq |\mathcal{B}_\ell| \frac{\ell}{2^{\ell+2}} - \varepsilon|X|n_1^{\ell-1} > \mu n_1^\ell \frac{\ell}{2^{\ell+2}} - \varepsilon n_1^\ell \geq 20\mu_{\ell-1}n_1^\ell. \quad (55)$$

Now, for each  $v \in X_2$  we count the number of  $\ell$ -sets  $B \in \mathcal{B}'_\ell$  containing  $v$ . We delete all vertices  $v$  from  $X_2$  for which this number is less than  $|\mathcal{B}'_\ell|/(10n_1)$  and call the resulting set  $X'$ . Observe that the definition of  $\mathcal{B}'_\ell$  implies that all vertices  $x$  in  $X'$  satisfy  $|N_Y(x)| \geq (d - \varepsilon)pn_2$ . Because  $\mathcal{B}'_\ell$  contains only 1-crossing  $\ell$ -sets we get

$$|\mathcal{B}'_\ell| \leq |X_2 \setminus X'| \frac{|\mathcal{B}'_\ell|}{10n_1} + |X'|n_1^{\ell-1} \leq \frac{|\mathcal{B}'_\ell|}{10} + |X'|n_1^{\ell-1}$$

and thus

$$|X'| \geq \frac{9}{10n_1^{\ell-1}} |\mathcal{B}'_\ell| \stackrel{(55)}{\geq} 10\mu_{\ell-1}n_1.$$

This together with Proposition 6 implies that the pairs  $(X', Y)$  and  $(Y, X_1)$  are  $(\varepsilon_{20}, d, p)$ -dense. In addition we have  $|X'|, |X_1| \geq \mu_{\ell-1}n_1 \geq \mu_{\ell-1}\xi p^{\Delta-1}n = \xi_{20}p^{\Delta-1}n$  and  $|Y| \geq \xi p^{\Delta-\ell}n \geq \xi_{20}p^{\Delta-2}n$ . Because  $\Gamma$  has property  $D(\varepsilon'_{20}, \mu_{20}, \varepsilon_{20}, \xi_{20}, p(n))$  we conclude for the tripartite graph  $G[X' \dot{\cup} Y \dot{\cup} X_1]$  that there are at least  $|X'| - \mu_{20}|X'| \geq 1$  vertices  $x$  in  $X'$  such that  $(N_Y(x), X_1)$  is  $(\varepsilon'_{20}, d, p)$ -dense. Let  $x^* \in X'$  be one of these vertices and set  $Y' := N_Y(x^*)$ . Thus  $(Y', X_1)$  is  $(\varepsilon'_{20}, d, p)$ -dense and since  $X'$  only contains vertices with a large neighbourhood in  $Y$  we have  $|Y'| \geq (d - \varepsilon)pn_2$ . Furthermore, let  $\mathcal{B}'_\ell(x^*)$  be the family of  $\ell$ -sets in  $\mathcal{B}'_\ell$  that contain  $x^*$ . Then  $\mathcal{B}'_\ell(x^*)$  contains  $\ell$ -sets with  $\ell - 1$  vertices in  $X_1$  and with one vertex, the vertex  $x^*$ , in  $X_2$  because  $\mathcal{B}'_\ell$  contains only 1-crossing  $\ell$ -sets. By definition of  $X'$  and because  $x^* \in X'$  we have

$$|\mathcal{B}'_\ell(x^*)| \geq |\mathcal{B}'_\ell|/(10n_1) \stackrel{(55)}{\geq} 2\mu_{\ell-1}n_1^{\ell-1}. \quad (56)$$

For  $B \in \mathcal{B}'_\ell(x^*)$  let  $\Pi_{\ell-1}(B)$  be the projection of  $B$  to  $X_1$ . This implies that  $\Pi_{\ell-1}(B)$  is an  $(\ell - 1)$ -set in  $X_1$ . In addition  $N_{Y'}(\Pi_{\ell-1}(B)) = N_Y(B)$  by definition of  $Y'$  and hence  $\Pi_{\ell-1}(B)$  has less than  $(d - \varepsilon')^\ell p^\ell n_2$  joint neighbours in  $Y'$  because  $B \in \mathcal{B}'_\ell(x^*) \subseteq \mathcal{B}_\ell$ . Accordingly the family  $\mathcal{B}_{\ell-1}$  of all projections  $\Pi_{\ell-1}(B)$  with  $B \in \mathcal{B}'_\ell(x^*)$  is a family of size  $|\mathcal{B}'_\ell(x^*)|$  and contains only  $(\ell - 1)$ -sets  $B'$  with

$$|N_{Y'}(B')| \leq (d - \varepsilon')^\ell p^\ell n_2 \leq (d - \varepsilon')^{\ell-1} p^{\ell-1} |Y'| = (d - \varepsilon'_{\ell-1})^{\ell-1} p^{\ell-1} |Y'|.$$

This means  $\mathcal{B}_{\ell-1} \subseteq \text{bad}_{\varepsilon'_{\ell-1}, d, p}^{G, \ell-1}(X_1, Y')$ . Recall that  $(X_1, Y')$  is  $(\varepsilon'_{20}, d, p)$ -dense by the choice of  $x^*$ . Because  $|X| = n_1 \geq \xi p^{\Delta-1}n$  and

$$|Y'| \geq (d - \varepsilon)pn_2 \geq (d - \varepsilon)p \cdot \xi p^{\Delta-\ell}n = \xi_{\ell-1}p^{\Delta-(\ell-1)}n$$

we can appeal to  $P_{\ell-1}(\varepsilon'_{\ell-1}, \mu_{\ell-1}, \varepsilon_{\ell-1}, \xi_{\ell-1}, p(n))$  and conclude that

$$|\mathcal{B}'_\ell(x^*)| = |\mathcal{B}_{\ell-1}| \leq |\text{bad}_{\varepsilon'_{\ell-1}, d, p}^{G, \ell-1}(X, Y')| \leq \mu_{\ell-1}n_1^{\ell-1},$$

contradicting (56).

Because  $G$  was arbitrary this shows that  $\Gamma$  has property  $P_\ell(\varepsilon', \mu, \varepsilon, \xi, p(n))$ . Thus  $(\mathcal{P}_\ell)$  holds, which finishes the proof of the inductive step.  $\square$

**B.3. Proof of Lemma 21.** In this section we provide the proof of Lemma 21 which examines the inheritance of  $p$ -density to neighbourhoods of  $\Delta$ -sets. For this purpose we will first establish a version of this lemma, Lemma 38 below, which only considers  $\Delta$ -sets that are crossing in a given vertex partition.

We need some definitions. Let  $G = (V, E)$  be a graph,  $X$  be a subset of its vertices, and  $X = X_1 \dot{\cup} \dots \dot{\cup} X_T$  be a partition of  $X$ . Then, for integers  $\ell, T > 0$ , we say that an  $\ell$ -set  $B \subseteq X$  is *crossing* in  $X_1 \dot{\cup} \dots \dot{\cup} X_T$  if there are indices  $0 < i_1 < \dots < i_\ell < T$  such that  $B$  contains exactly

one element in  $X_{i_j}$  for each  $j \in [\ell]$ . In this case we also write  $B \in X_{i_1} \times \dots \times X_{i_\ell}$  (hence identifying crossing  $\ell$ -sets with  $\ell$ -tuples).

Now let  $p, \varepsilon, d$  be positive reals, and  $Y, Z \subseteq V$  be vertex sets such that  $X, Y$ , and  $Z$  are mutually disjoint. Define

$$\text{bad}_{\varepsilon, d, p}^{G, \ell}(X_1, \dots, X_T; Y, Z)$$

to be the family of all those crossing  $\ell$ -sets  $B$  in  $X_1 \dot{\cup} \dots \dot{\cup} X_T$  that either satisfy  $|N_Y^\cap(B)| < (d - \varepsilon)^\ell p^\ell |Y|$  or have the property that  $(N_Y^\cap(B), Z)$  is not  $(\varepsilon, d, p)$ -dense in  $G$ . Further, let

$$\text{Bad}_{\varepsilon, d, p}^{G, \ell}(X_1, \dots, X_T; Y, Z)$$

be the family of crossing  $\ell$ -sets  $B$  in  $X_1 \dot{\cup} \dots \dot{\cup} X_T$  that contain an  $\ell'$ -set  $B' \subseteq B$  with  $\ell' > 0$  such that  $B' \in \text{bad}_{\varepsilon, d, p}^{G, \ell'}(X_1, \dots, X_T; Y, Z)$ .

**Lemma 38.** *For all integers  $\ell, \Delta > 0$  and positive reals  $d_0, \varepsilon'$ , and  $\mu$  there is  $\varepsilon$  such that for all  $\xi > 0$  there is  $c > 1$  such that if  $p > c(\frac{\log n}{n})^{1/\Delta}$ , then the following holds a.a.s. for  $\Gamma = \mathcal{G}_{n, p}$ . For  $n_1, n_3 \geq \xi p^{\Delta-1}n$  and  $n_2 \geq \xi p^{\Delta-\ell-1}n$  let  $G = (X \dot{\cup} Y \dot{\cup} Z, E)$  be any tripartite subgraph of  $\Gamma$  with  $|X| = n_1, |Y| = n_2$ , and  $|Z| = n_3$ . Assume further that  $X = X_1 \dot{\cup} \dots \dot{\cup} X_\ell$  with  $|X_i| \geq \lfloor \frac{n_1}{\ell} \rfloor$  and that  $(X, Y)$  and  $(Y, Z)$  are  $(\varepsilon, d, p)$ -dense pairs with  $d \geq d_0$ . Then*

$$|\text{bad}_{\varepsilon', d, p}^{G, \ell}(X_1, \dots, X_\ell; Y, Z)| \leq \mu n_1^\ell.$$

*Proof.* Let  $\Delta$  and  $d_0$  be given. For a fixed  $n$ -vertex graph  $\Gamma$ , a fixed integer  $\ell$  and fixed positive reals  $\varepsilon', \mu, \varepsilon, \xi$ , and a function  $p = p(n)$  we say that we say that a graph  $\Gamma$  on  $n$  vertices has property  $\mathcal{P}_\ell(\varepsilon', \mu, \varepsilon, \xi, p(n))$  if  $\Gamma$  has the property stated in the lemma for these parameters and for  $\Delta$  and  $d_0$ , that is, whenever  $G = (X \dot{\cup} Y \dot{\cup} Z, E)$  is a tripartite subgraph of  $\Gamma$  with the required properties, then  $G$  satisfies the conclusion of the lemma. For any fixed  $\ell > 0$ , we denote by  $(\mathcal{P}_\ell)$  the following statement.

$(\mathcal{P}_\ell)$  *For all  $\varepsilon', \mu > 0$  there is  $\varepsilon$  such that for all  $\xi > 0$  there is  $c > 1$  such that a random graph  $\Gamma = \mathcal{G}_{n, p}$  with  $p > c(\frac{\log n}{n})^{1/\Delta}$  has property  $\mathcal{P}_\ell(\varepsilon', \mu, \varepsilon, \xi, p(n))$  with probability  $1 - o(1)$ .*

We prove that  $(\mathcal{P}_\ell)$  holds for every fixed  $\ell > 0$  by induction on  $\ell$ .

The case  $\ell = 1$  is an easy consequence of Lemma 20 and Proposition 7. Indeed, let  $\varepsilon'$  and  $\mu$  be arbitrary, let  $\varepsilon_{20}$  be as asserted by Lemma 20 for  $\Delta, d_0, \varepsilon'$ , and  $\mu/2$  and fix  $\varepsilon := \min\{\varepsilon_{20}, \varepsilon', \mu/2\}$ . Let  $\xi$  be arbitrary and pass it on to Lemma 20 for obtaining  $c$ . Now, let  $\Gamma = \mathcal{G}_{n, p}$  be a random graph. Then, by the choice of parameters, Lemma 20 asserts that the graph  $\Gamma$  has the following property with probability  $1 - o(1)$ . Let  $G = (X \dot{\cup} Y \dot{\cup} Z, E)$  be any subgraph with  $X = X_1$  and  $|X| = n_1, |Y| = n_2$ , and  $|Z| = n_3$ , where  $n_1, n_3 \geq \xi p^{\Delta-1}n$  and  $n_2 \geq \xi p^{\Delta-2}n$ , and  $(X, Y)$  and  $(Y, Z)$  are  $(\varepsilon, d, p)$ -dense pairs. Then there are at most  $\frac{\mu}{2}n_1$  vertices  $x \in X$  such that  $(N(x) \cap Y, Z)$  is not an  $(\varepsilon', d, p)$ -dense pair in  $G$ . Because  $\varepsilon \leq \mu/2$ , Proposition 7 asserts that in every such  $G$  there are at most  $\frac{\mu}{2}n_1$  vertices  $x \in X$  with  $|N_Y(x)| < (d - \varepsilon')p|Y|$ . This implies that

$$|\text{bad}_{\varepsilon', d, p}^{G, 1}(X_1; Y, Z)| \leq \mu n_1$$

holds with probability  $1 - o(1)$  for all such subgraphs  $G$  of the random graph  $\Gamma$ . Accordingly we get  $(\mathcal{P}_1)$ .

For the inductive step assume that  $(\mathcal{P}_{\ell-1})$  and  $(\mathcal{P}_1)$  hold. We will show that this implies  $(\mathcal{P}_\ell)$ . Again, let  $\varepsilon'$  and  $\mu$  be arbitrary positive constants. Let  $\varepsilon_1$  be as promised in the statement  $(\mathcal{P}_1)$  for parameters  $\varepsilon'_1 := \varepsilon'$  and  $\mu_1 := \mu/2$ . Set  $\varepsilon'_{\ell-1} := \min\{\varepsilon_1, \varepsilon', \frac{\mu}{4}\}$ , and let  $\varepsilon_{\ell-1}$  be given by  $(\mathcal{P}_{\ell-1})$  for parameters  $\varepsilon'_{\ell-1}$  and  $\mu_{\ell-1} := \frac{\mu}{4}$ . We define  $\varepsilon := \varepsilon_{\ell-1}/(\ell + 1)$ . Next, let  $\xi$  be an arbitrary parameter and choose

$$\xi_1 := \min\{\xi/(\ell + 1), (d_0 - \varepsilon'_{\ell-1})^{\ell-1}\xi\} \quad \text{and} \quad \xi_{\ell-1} := \xi/(\ell + 1). \quad (57)$$

Finally, let  $c_1$  and  $c_{\ell-1}$  be given by  $(\mathcal{P}_1)$  and  $(\mathcal{P}_{\ell-1})$ , respectively, for the previously specified parameters together with  $\xi_1$  and  $\xi_{\ell-1}$ . Set  $c := \max\{c_1, c_{\ell-1}\}$ . We will prove that with this choice of  $\varepsilon$  and  $c$  the statement in  $(\mathcal{P}_\ell)$  holds for the input parameters  $\varepsilon', \mu$ , and  $\xi$ . For this purpose let  $\Gamma = \mathcal{G}_{n, p}$  be a random graph. By  $(\mathcal{P}_1)$  and  $(\mathcal{P}_{\ell-1})$  and the choice of the parameters the graph  $\Gamma$  has properties  $\mathcal{P}_1(\varepsilon'_1, \mu_1, \varepsilon_1, \xi_1, p(n))$  and  $\mathcal{P}_{\ell-1}(\varepsilon'_{\ell-1}, \mu_{\ell-1}, \varepsilon_{\ell-1}, \xi_{\ell-1}, p(n))$  with probability

$1 - o(1)$ . We will show that a graph  $\Gamma$  with these properties also satisfies  $P_\ell(\varepsilon', \mu, \varepsilon, \xi, p(n))$ . Let  $G = (X \dot{\cup} Y \dot{\cup} Z, E)$  be an arbitrary subgraph of such a  $\Gamma$  where  $X = X_1 \dot{\cup} \dots \dot{\cup} X_\ell$ ,  $|X| = n_1$ ,  $|Y| = n_2$ ,  $|Z| = n_3$ , with  $n_1, n_3 \geq \xi p^{\Delta-1} n$ ,  $n_2 \geq \xi p^{\Delta-\ell-1} n$ , and  $|X_i| \geq \lfloor \frac{n_1}{\ell} \rfloor$ , and assume that  $(X, Y)$  and  $(Y, Z)$  are  $(\varepsilon, d, p)$ -dense pairs for  $d \geq d_0$ .

We would like to bound  $\mathcal{B}_\ell := \text{bad}_{\varepsilon', d, p}^{G, \ell}(X_1, \dots, X_\ell; Y, Z)$ . For this purpose let  $B'$  be a fixed  $(\ell - 1)$ -set and define

$$\mathcal{B}_{\ell-1} := \text{bad}_{\varepsilon'_{\ell-1}, d, p}^{G, \ell-1}(X_1, \dots, X_{\ell-1}; Y, Z) \cup \text{bad}_{\varepsilon'_{\ell-1}, d, p}^{G, \ell-1}(X_1, \dots, X_{\ell-1}; Y, X_\ell) \quad (58a)$$

$$\mathcal{B}_1(B') := \text{bad}_{\varepsilon', d, p}^{G, 1}(X_\ell; N_Y^\cap(B'), Z). \quad (58b)$$

For an  $\ell$ -set  $B \in X_1 \times \dots \times X_\ell$  let further  $\Pi_{\ell-1}(B)$  denote the  $(\ell - 1)$ -set that is the projection of  $B$  to  $X_1 \times \dots \times X_{\ell-1}$  and let  $\Pi_\ell(B)$  be the vertex that is the projection of  $B$  to  $X_\ell$ . Now, consider an  $\ell$ -set  $B$  that is contained in  $B \in \mathcal{B}_\ell$  but is such that  $B' := \Pi_{\ell-1}(B) \notin \mathcal{B}_{\ell-1}$ . Let  $v = \Pi_\ell(B) \in X_\ell$  and  $Y' := N_Y^\cap(B')$ . We will show that then  $v \in \mathcal{B}_1(B')$ . Indeed, since  $B' \notin \mathcal{B}_{\ell-1}$  it follows from (58a) that

$$B' \notin \text{bad}_{\varepsilon'_{\ell-1}, d, p}^{G, \ell-1}(X_1, \dots, X_{\ell-1}; Y, Z)$$

and thus  $|Y'| \geq (d - \varepsilon'_{\ell-1})^{\ell-1} p^{\ell-1} n_2$ . As  $N_Y^\cap(B) = N_Y^\cap(v)$  we conclude that

$$v \in \text{bad}_{\varepsilon', d, p}^{G, 1}(X_\ell; Y', Z) = \mathcal{B}_1(B')$$

by (58b) because otherwise  $(N_Y^\cap(B), Z)$  was  $(\varepsilon', d, p)$ -dense and we had

$$|N_Y^\cap(B)| \geq (d - \varepsilon') p |Y'| \geq (d - \varepsilon') p \cdot (d - \varepsilon'_{\ell-1})^{\ell-1} p^{\ell-1} n_2 \geq (d - \varepsilon')^\ell p^\ell n_2,$$

which contradicts  $B \in \mathcal{B}_\ell$ . Summarizing, we have

$$\begin{aligned} \mathcal{B}_\ell &= \{B \in \mathcal{B}_\ell : \Pi_{\ell-1}(B) \in \mathcal{B}_{\ell-1}\} \cup \{B \in \mathcal{B}_\ell : \Pi_{\ell-1}(B) \notin \mathcal{B}_{\ell-1}\} \\ &\subseteq (\mathcal{B}_{\ell-1} \times X_\ell) \cup \bigcup_{B' \notin \mathcal{B}_{\ell-1}} \{B'\} \times \mathcal{B}_1(B'). \end{aligned} \quad (59)$$

For bounding  $\mathcal{B}_\ell$  we will thus estimate the sizes of  $\mathcal{B}_{\ell-1}$  and  $\mathcal{B}_1(B')$  for  $B' \notin \mathcal{B}_{\ell-1}$ . Let  $X' := X_1 \dot{\cup} \dots \dot{\cup} X_{\ell-1}$ . Since  $(X, Y)$  is  $(\varepsilon, d, p)$ -dense we conclude from Proposition 6 that  $(X', Y)$  and  $(X_\ell, Y)$  are  $(\varepsilon_{\ell-1}, d, p)$ -dense pairs since  $\varepsilon(\ell + 1) \leq \varepsilon_{\ell-1}$ . Further, by the choice of  $\xi_{\ell-1}$  we get  $|X'|, |X_\ell| \geq n_1/(\ell + 1) \geq \xi_{\ell-1} p^{\Delta-1} n$  since  $n_1 \geq \xi p^{\Delta-1} n$  by assumption. Thus we can use the fact that  $\Gamma$  has property  $P_{\ell-1}(\varepsilon'_{\ell-1}, \mu_{\ell-1}, \varepsilon_{\ell-1}, \xi_{\ell-1}, p(n))$  once on the tripartite subgraph induced on  $X' \dot{\cup} Y \dot{\cup} Z$  in  $G$  and once on the tripartite subgraph induced on  $X' \dot{\cup} Y \dot{\cup} X_\ell$  in  $G$  and infer that

$$|\mathcal{B}_{\ell-1}| \leq 2 \cdot \mu_{\ell-1} n_1^{\ell-2} = \frac{\mu}{2} n_1^{\ell-2}. \quad (60)$$

For estimating  $|\mathcal{B}_1(B')|$  for  $B' \notin \mathcal{B}_{\ell-1}$  let  $Y' := N_Y^\cap(B')$ . Observe that this implies that  $(Y', Z)$  and  $(X_\ell, Y')$  are  $(\varepsilon_1, d, p)$ -dense pairs because  $\varepsilon'_{\ell-1} \leq \varepsilon_1$ , and that

$$|Y'| \geq (d - \varepsilon'_{\ell-1})^{\ell-1} p^{\ell-1} n_2 \geq (d - \varepsilon'_{\ell-1})^{\ell-1} p^{\ell-1} \cdot \xi p^{\Delta-\ell-1} n \stackrel{(57)}{\geq} \xi_1 p^{\Delta-1} n.$$

By (57)  $|X_\ell|, |Z| \geq \xi p^{\Delta-1} n/(\ell + 1) \geq \xi_1 p^{\Delta-1} n$ . As  $\Gamma$  satisfies  $P_1(\varepsilon'_1, \mu_1, \varepsilon_1, \xi_1, p(n))$  we conclude that

$$|\mathcal{B}_1(B')| \stackrel{(58b)}{=} |\text{bad}_{\varepsilon', d, p}^{G, 1}(X_\ell; Y', Z)| \leq \mu_1 n_1 \leq \frac{\mu}{2} n_1. \quad (61)$$

In view of (59), combining (60) and (61) gives

$$\left| \text{bad}_{\varepsilon', d, p}^{G, \ell}(X_1, \dots, X_\ell; Y, Z) \right| = |\mathcal{B}_\ell| \leq \frac{\mu}{2} n_1^{\ell-1} \cdot n_1 + n_1^{\ell-1} \cdot \frac{\mu}{2} n_1 = \mu n_1^\ell.$$

Because  $G$  was arbitrary this shows that  $\Gamma$  has property  $P_\ell(\varepsilon', \mu, \varepsilon, \xi, p(n))$ . Thus  $(\mathcal{P}_\ell)$  holds which finishes the proof of the inductive step.  $\square$

In the proof of Lemma 21 we now first partition the vertex set  $X$ , in which we count bad  $\ell$ -sets, arbitrarily into  $T$  vertex sets of equal size. Lemma 38 then implies that for all  $\ell' \in [\ell]$  there are not many bad  $\ell'$ -sets that are crossing in this partition. It follows that only few  $\ell$ -sets in  $X$  contain a bad  $\ell'$ -set for some  $\ell' \in [\ell]$  (recall that in Definition 17 for  $\text{Bad}_{\varepsilon, d, p}^{G, \ell}(X, Y, Z)$  such



$\ell'$ -sets are considered). Moreover, if  $T$  is sufficiently large then the number of non-crossing  $\ell$ -sets is negligible. Hence we obtain that there are few bad sets in total.

*Proof of Lemma 21.* Given  $\Delta, \ell, d_0, \varepsilon'$  and  $\mu$  let  $T$  be such that  $\mu T \geq 2$ , fix  $\mu_{38} := \frac{1}{2}\mu/(\ell T^\ell)$ . For  $j \in [\ell]$  let  $\varepsilon_j$  be given by Lemma 38 with  $\ell$  replaced by  $j$  and for  $\Delta, d_0, \varepsilon'$ , and  $\mu_{38}$  and set  $\varepsilon_{38} := \min_{j \in [\ell]} \varepsilon_j$ . Define  $\varepsilon := \varepsilon_{38}/(T+1)$ . Now, in Lemma 21 let  $\xi$  be given by the adversary for this  $\varepsilon$ . Set  $\xi_{38} := \xi/(T+1)$ , and let  $c$  be given by Lemma 38 for this  $\xi_{38}$ .

Let  $\Gamma = \mathcal{G}_{n,p}$  with  $p \geq c(\frac{\log n}{n})^{1/\Delta}$ . Then a.a.s. the graph  $\Gamma$  satisfies the statement in Lemma 38 for parameters  $j \in [\ell]$ ,  $\Delta, d_0, \varepsilon', \mu_{38}$ , and  $\xi_{38}$ . Assume that  $\Gamma$  has this property for all  $j \in [\ell]$ . We will show that it then also satisfies the statement in Lemma 21.

Indeed, let  $G$  and  $X, Y, Z$  be arbitrary with the properties as required in Lemma 21. Let  $X = X_1 \dot{\cup} \dots \dot{\cup} X_T$  be an arbitrary partition of  $X$  with  $|X_i| \geq \lfloor \frac{n_1}{T} \rfloor$ . We will first show that there are not many bad crossing  $\ell$ -sets with respect to this partition, i.e., we will bound the size of  $\text{Bad}_{\varepsilon', d, p}^{G, \ell}(X_1, \dots, X_T; Y, Z)$ . By definition

$$|\text{Bad}_{\varepsilon', d, p}^{G, \ell}(X_1, \dots, X_T; Y, Z)| \leq \sum_{j \in [\ell]} |\text{bad}_{\varepsilon', d, p}^{G, j}(X_1, \dots, X_T; Y, Z)| \cdot n_1^{\ell-j}.$$

Now, fix  $j \in [\ell]$  and an index set  $\{i_1, \dots, i_j\} \in \binom{[T]}{j}$  and consider the induced tripartite subgraph  $G' = (X' \dot{\cup} Y \dot{\cup} Z, E')$  of  $G$  with  $X' = X_{i_1} \dot{\cup} \dots \dot{\cup} X_{i_j}$ . Observe that  $|Y| \geq \xi_{38} p^{\Delta-j-1} n$ ,  $|Z| \geq \xi_{38} p^{\Delta-1} n$ , and  $n'_1 := |X'| \geq j \lfloor n_1/T \rfloor \geq \xi_{38} p^{\Delta-1} n$ . By definition  $\varepsilon(T+1)/j \leq \varepsilon_{38} \leq \varepsilon_j$  and so by Proposition 6 the pair  $(X', Y)$  is  $(\varepsilon_j, d, p)$ -dense. Thus, because  $\Gamma$  satisfies the statement in Lemma 38 for parameters  $j, \Delta, d, \varepsilon', \mu_{38}$ , and  $\xi_{38}$  we have that  $G'$  satisfies

$$|\text{bad}_{\varepsilon', d, p}^{G', j}(X_{i_1}, \dots, X_{i_j}; Y, Z)| \leq \mu_{38} (n'_1)^j.$$

As there are  $\binom{T}{j}$  choices for the index set  $\{i_1, \dots, i_j\}$  this implies

$$|\text{bad}_{\varepsilon', d, p}^{G, j}(X_1, \dots, X_T; Y, Z)| \leq \binom{T}{j} \mu_{38} (n'_1)^j \leq T^j \mu_{38} n_1^j,$$

and thus

$$|\text{Bad}_{\varepsilon', d, p}^{G, \ell}(X_1, \dots, X_T; Y, Z)| \leq \sum_{j \in [\ell]} T^j \mu_{38} n_1^j \cdot n_1^{\ell-j} \leq \frac{1}{2} \mu n_1^\ell.$$

The number of  $\ell$ -sets in  $X$  that are not crossing with respect to the partition  $X = X_1 \dot{\cup} \dots \dot{\cup} X_T$  is at most  $T \binom{n_1/T}{\ell-2} \leq \frac{1}{T} n_1^\ell \leq \frac{1}{2} \mu n_1^\ell$  and so we get  $|\text{Bad}_{\varepsilon', d, p}^{G, \ell}(X, Y, Z)| \leq \mu n_1^\ell$ .  $\square$

INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, RUA DO MATÃO 1010, 05508-090 SÃO PAULO, BRAZIL

*E-mail address:* julia@ime.usp.br

INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, RUA DO MATÃO 1010, 05508-090 SÃO PAULO, BRAZIL

*E-mail address:* yoshi@ime.usp.br

ZENTRUM MATHEMATIK, TECHNISCHE UNIVERSITÄT MÜNCHEN, BOLTZMANNSTRASSE 3, D-85747 GARCHING BEI MÜNCHEN, GERMANY

*E-mail address:* taraz@ma.tum.de